

April 25, 2017

REDUCIBILITY OF QUANTUM HARMONIC OSCILLATOR ON $\mathbb{R}^d$ WITH DIFFERENTIAL AND QUASI-PERIODIC IN TIME POTENTIAL

ZHENGUO LIANG AND ZHIGUO WANG

ABSTRACT. We improve the results by Grébert and Paturel in [22] and prove that a linear Schrödinger equation on $\mathbb{R}^d$ with harmonic potential $|x|^2$ and small t -quasiperiodic potential as

$$iu_t - \Delta u + |x|^2 u + \varepsilon V(\omega t, x)u = 0, \quad (t, x) \in \mathbb{R} \times \mathbb{R}^d$$

reduces to an autonomous system for most values of the frequency vector $\omega \in \mathbb{R}^n$. The new point is that the potential $V(\theta, \cdot)$ is only in $C^\beta(\mathbb{T}^n, \mathcal{H}^s(\mathbb{R}^d))$ with β large enough. As a consequence any solution of such a linear PDE is almost periodic in time and remains bounded in some suitable Sobolev norms.

KEYWORDS. quantum harmonic oscillator, finitely differentiable, pure-point spectrum, KAM, reducibility

MATHEMATICS SUBJECT CLASSIFICATION NUMBERS. 35P05, 37K55, 81Q15

1. INTRODUCTION

1.1. State of Reducibility Problem and Main Results. We consider the following nonautonomous linear equation in $\mathbb{R}^d$

$$iu_t - \Delta u + |x|^2 u + \varepsilon V(\omega t, x)u = 0, \quad u = u(t, x), \quad (t, x) \in \mathbb{R} \times \mathbb{R}^d. \quad (1.1)$$

Here $\varepsilon > 0$ is a small parameter and the frequency vector ω of forced oscillator is regarded as a parameter in $D_0 = [0, 2\pi]^n \subset \mathbb{R}^n$. The function V is a real multiplicative potential which is quasiperiodic in time. Namely, V is a continuous function of $(\theta, x) \in \mathbb{T}^n \times \mathbb{R}^d$. We assume $V(z, \cdot) \in C^\beta(\mathbb{R}^n, \mathcal{H}^s(\mathbb{R}^d))$ which will be denoted in the following (see Definition 1.1).

As the usual reducibility results we consider the previous equation as a linear non-autonomous equation in the complex Hilbert space $L^2(\mathbb{R}^d)$ and we prove that it reduces to an autonomous system for most values of the frequency vector ω .

Similar as Grébert and Paturel [22], we introduce some notations. Let $T := -\Delta + |x|^2 = -\Delta + x_1^2 + \cdots + x_d^2$ be the d -dimensional quantum harmonic oscillator. Its spectrum is the sum of d copies of odd integers, i.e., the spectrum of T equals to $\hat{\mathcal{E}} := \{d, d+2, d+4, \cdots\}$. For $j \in \hat{\mathcal{E}}$, we denote the associated eigenspace by E_j whose dimension is

$$d_j := \text{card}\{(i_1, \cdots, i_d) \in (2\mathbb{N}-1)^d \mid i_1 + \cdots + i_d = j\} \leq j^{d-1}.$$

We denote $\{\Phi_{j,l}, l = 1, 2, \cdots, d_j\}$, the basis of E_j obtained by d -tensor product of Hermite functions: $\Phi_{j,l} = \varphi_{i_1} \otimes \cdots \otimes \varphi_{i_d}$ for some choice of $i_1 + \cdots + i_d = j$. Then setting

$$\mathcal{E} := \{(j, l) \in \hat{\mathcal{E}} \times \mathbb{N} \mid l = 1, \cdots, d_j\}.$$

$(\Phi_a)_{a \in \mathcal{E}}$ is a basis of $L^2(\mathbb{R}^d)$ and denote $w_{j,l} := j$ for $(j, l) \in \mathcal{E}$. We have

$$T\Phi_a = w_a \Phi_a, \quad a \in \mathcal{E}. \quad (1.2)$$

The first author was partially supported by NSFC grants 11371097, 11571249; the second author was partially supported by NSFC grants 11571249, 11671192.

We define in $\mathcal{E}$ an equivalence relation $a \sim b \Leftrightarrow w_a = w_b$ and denote by $[a]$ the equivalence class associated to $a \in \mathcal{E}$. Note that $\text{card } [a] \leq w_a^{d-1}$.

Let $s \geq 0$ be an integer we define

$$\mathcal{H}^s := \{u \in \mathcal{H}^s(\mathbb{R}^d, \mathbb{C}) \mid x \mapsto x^{\alpha_1} \partial_x^{\alpha_2} u \in L^2(\mathbb{R}^d) \text{ for any } \alpha_1, \alpha_2 \in \mathbb{N}^d, 0 \leq |\alpha_1| + |\alpha_2| \leq s\}.$$

We need to point it out that $\mathcal{H}^s$ is the form domain of T^s and the domain of $T^{\frac{s}{2}}$ and this allows us to extend the definition of $\mathcal{H}^s$ to any nonnegative real values of s (see Delort [14]).

To a function $u \in \mathcal{H}^s$ we associate the sequence ξ of its Hermite coefficients by the formula $u(x) = \sum_{a \in \mathcal{E}} \xi_a \Phi_a(x)$. Then we define $\ell_s^2 := \{(\xi)_{a \in \mathcal{E}} \mid \sum_{a \in \mathcal{E}} w_a^s |\xi_a|^2 < \infty\}$, and for $s \geq 0$, $u \in \mathcal{H}^s \Leftrightarrow \xi \in \ell_s^2$. Then we endow both spaces with the norm $\|u\|_s = \|\xi\|_s = (\sum_{a \in \mathcal{E}} w_a^s |\xi_a|^2)^{\frac{1}{2}}$. If s is a nonnegative integer, we will use the fact that the norm on $\mathcal{H}^s$ are equivalently defined as $\|T^{\frac{s}{2}} f\|_{L^2(\mathbb{R}^d)}$ and $\sum_{0 \leq |\alpha_1| + |\alpha_2| \leq s} \|x^{\alpha_1} \partial_x^{\alpha_2} f\|_{L^2(\mathbb{R}^d)}$.

To introduce the main result we introduce some notations and definitions.

Definition 1.1. Assume that X is a complex Banach space with the norm $\|\cdot\|_X$. Let $\mathcal{C}^b(\mathbb{R}^n, X)$, $0 < b < 1$, be the space of Hölder continuous functions $f : \mathbb{R}^n \rightarrow X$ with the norm

$$\|f\|_{\mathcal{C}^b(\mathbb{R}^n, X)} := \sup_{0 < |z_1 - z_2| < 2\pi} \frac{\|f(z_1) - f(z_2)\|_X}{|z_1 - z_2|^b} + \sup_{z \in \mathbb{R}^n} \|f(z)\|_X.$$

If $b = 0$, then $\|f\|_{\mathcal{C}^b(\mathbb{R}^n, X)}$ denotes the sup-norm. For $\beta = [\beta] + b$ with $0 \leq b < 1$, we denote by $\mathcal{C}^\beta(\mathbb{R}^n, X)$ the space of functions $f : \mathbb{R}^n \rightarrow X$ with Hölder continuous partial derivatives and $\partial^\alpha f \in \mathcal{C}^b(\mathbb{R}^n, X_\alpha)$ for all multi-indices $\alpha = (\alpha_1, \dots, \alpha_n) \in \mathbb{N}^n$, where $|\alpha| := |\alpha_1| + \dots + |\alpha_n| \leq \beta$ and $X_\alpha = \mathfrak{L}(\prod_{i=1}^{|\alpha|} Y_i, X)$ with the standard norm and $Y_i := \mathbb{R}^n$, $i = 1, \dots, |\alpha|$. We define the norm $\|f\|_{\mathcal{C}^\beta(\mathbb{R}^n, X)} := \sum_{|\alpha| \leq \beta} \|\partial^\alpha f\|_{\mathcal{C}^b(\mathbb{R}^n, X_\alpha)}$. If a function f has a finite norm $\|f\|_{\mathcal{C}^\beta(\mathbb{R}^n, X)}$, then we call $f \in \mathcal{C}^\beta(\mathbb{R}^n, X)$.

Denote by $\mathcal{C}^\beta(\mathbb{T}^n, X)$ the space of all functions $f \in \mathcal{C}^\beta(\mathbb{R}^n, X)$ that are of period 2π in all variables. We define $\|f\|_{\mathcal{C}^\beta(\mathbb{T}^n, X)} := \|f\|_{\mathcal{C}^\beta(\mathbb{R}^n, X)}$.

Definition 1.2. A real potential $V : \mathbb{T}^n \times \mathbb{R}^d \ni (\theta, x) \mapsto V(\theta, x)$ is called (s, β) -admissible if $V(\theta, x) \in \mathcal{C}^\beta(\mathbb{T}^n, \mathcal{H}^s(\mathbb{R}^d))$ with a finite norm, namely, $\|V(\theta, \cdot)\|_{\mathcal{C}^\beta(\mathbb{T}^n, \mathcal{H}^s(\mathbb{R}^d))} \leq C$, where

$$\begin{cases} s \geq 0, & d = 1, \\ s > 2(d-2), & d \geq 2, \end{cases}$$

and the constant C depends on s, β, n and d .

Set $\gamma_1 = n + d + 2$, $\gamma_2 = \frac{\alpha}{4+d+2\alpha}$ (depending only on s and d) and α given by (3.1), we have

Theorem 1.3. Assume that the potential $V : \mathbb{T}^n \times \mathbb{R}^d \ni (\theta, x) \mapsto \mathbb{R}$ is (s, β) -admissible. There exists δ satisfying $0 < \delta < \frac{\gamma_2}{24}$ and $\varepsilon_*(\beta, n, s, d, \delta) > 0$, if $0 < \varepsilon < \varepsilon_*$, $\beta > \max\{9(2 + \frac{d}{\alpha}) - \frac{\gamma_1}{\gamma_2 - 24\delta}, 9n, 12(d+1)\}$, then there exists $D_\varepsilon \subset D_0$ with $\text{Meas}(D_0 \setminus D_\varepsilon) \leq c(\beta, n, d, s, \delta) \varepsilon^{\frac{3\delta}{2+\frac{d}{\alpha}}}$, such that for all $\omega \in D_\varepsilon$ the linear Schrödinger equation (1.1) reduces to a linear autonomous equation in the space $\mathcal{H}^{s'}$ with $1 \leq s' \leq \max\{s, 1\}$.

More precisely, for $\omega \in D_\varepsilon$, there exist a linear isomorphism $\Psi_\omega^\infty(\theta) \in \mathfrak{L}(\mathcal{H}^{s'})$ for $0 \leq s' \leq s$, unitary on $L^2(\mathbb{R}^d)$, where $\Psi_\omega^\infty(\theta) \in \mathcal{C}^\mu(\mathbb{T}^n, \mathfrak{L}(\mathcal{H}^{s'}))$ for $0 \leq s' \leq s$ with $\mu \notin \mathbb{Z}$ and $\mu \leq \frac{2}{9}\beta$, and a bounded Hermitian operator $W = W_{\omega, \varepsilon} \in \mathfrak{L}(\mathcal{H}^{s'})$ such that $t \mapsto u(t, \cdot) \in \mathcal{H}^{s'}$ with $1 \leq s' \leq \max\{s, 1\}$ satisfies (1.1) if and only if $t \mapsto v(t, \cdot) = \Psi_\omega^\infty(\omega t)u(t, \cdot)$ satisfies the autonomous equation

$$i\partial_t v - \Delta v + |x|^2 v + \varepsilon W(v) = 0.$$

Furthermore, for $0 \leq s' \leq s$,

$$\|\Psi_\omega^\infty(\theta) - id\|_{\mathcal{C}^\mu(\mathbb{T}^n, \mathfrak{L}(\mathcal{H}^{s'}, \mathcal{H}^{s'+2\alpha}))} \leq C\varepsilon^{\frac{3}{2\beta}(\frac{2}{3}\beta - \mu)}, \quad (\theta, \omega) \in \mathbb{T}^n \times D_\varepsilon.$$

On the other hand, the infinite matrix $(W_a^b)_{a,b \in \mathcal{E}}$ of the operator W written in the Hermite basis ($W_a^b = \int_{\mathbb{R}^d} \Phi_a W(\Phi_b) dx$) is block diagonal, i.e. $W_a^b = 0$ if $w_a \neq w_b$. Denote $(\langle V \rangle_a^b)_{a,b \in \mathcal{E}}$ be the corresponding infinite matrix of the operator $\langle V \rangle(x) = \frac{1}{(2\pi)^n} \int_{\mathbb{T}^n} V(\theta, x) d\theta$ written in the Hermite basis, we have

$$\|(W_a^b)_{a,b \in \mathcal{E}} - \Pi((\langle V \rangle_a^b)_{a,b \in \mathcal{E}})\|_{\mathfrak{L}(\ell_{s'}^2)} \leq c\varepsilon^{\frac{1}{2}}$$

for $0 \leq s' \leq s$, where Π is the projection on the diagonal blocks.

Remark 1.4. Comparing with Theorem 1.2 in [22] we prove the reducibility theorem in the space $\mathcal{H}^{s'}$ with $1 \leq s' \leq \max\{s, 1\}$, not in the energy space $\mathcal{H}^1$. See Lemma 4.28 for details.

As a consequence of Theorem 1.3, we prove the following corollary concerning the solutions of (1.1).

Corollary 1.5. Assume all the assumptions in Theorem 1.3 hold. Let $1 \leq s' \leq \max\{s, 1\}$ and let $u_0 \in \mathcal{H}^{s'}$, then there exists $\varepsilon_* > 0$ such that for $0 < \varepsilon < \varepsilon_*$ and $\omega \in D_\varepsilon$ (in Theorem 1.3), there exists a unique solution $u \in \mathcal{C}(\mathbb{R}, \mathcal{H}^{s'})$ of (1.1) such that $u(0) = u_0$. Moreover, u is almost periodic in time and satisfies

$$(1 - c\varepsilon)\|u_0\|_{\mathcal{H}^{s'}} \leq \|u(t)\|_{\mathcal{H}^{s'}} \leq (1 + c\varepsilon)\|u_0\|_{\mathcal{H}^{s'}}, \quad \forall t \in \mathbb{R}$$

with some $c = c(s', s, d)$.

Consider on $L^2(\mathbb{T}^n) \otimes L^2(\mathbb{R}^d)$ the Floquet Hamiltonian operator

$$K := -i \sum_{k=1}^n \omega_k \frac{\partial}{\partial \theta_k} - \Delta + |x|^2 + \varepsilon V(\theta, x),$$

we have

Corollary 1.6. Assume all the assumptions in Theorem 1.3 hold. Then there exists $\varepsilon_* > 0$ such that for $0 < \varepsilon < \varepsilon_*$ and $\omega \in D_\varepsilon$, the spectrum of the Floquet operator K is pure point.

1.2. Related results. The equations (1.1) can be generalized into a time-dependent Schrödinger equation

$$i\partial_t \zeta(t) = (\mathcal{A} + \varepsilon \mathcal{B}(\omega t)) \zeta(t), \quad (1.3)$$

where $\mathcal{A}$ is a positive self-adjoint operator on a separable Hilbert space $\mathcal{H}$ and the perturbation $\mathcal{B}$ is an operator-valued function from $\mathbb{T}^n$ into the space of symmetric operators on $\mathcal{H}$. Our aim is to show that for sufficiently small ε , and for ω belonging to a set of large measure, there exists a unitary transformation which conjugates Eq. (1.3) to a time independent equation. If this is true, we will call Eq. (1.3) is *reducible*. From the reducibility of Eq. (1.3) and relative properties of the transformation we can easily prove the boundedness of the Sobolev norms and pure point spectrum of the relative Floquet operator, which is defined by

$$K_F := -i\omega \cdot \partial_\theta + \mathcal{A} + \varepsilon \mathcal{B}(\theta) \quad \text{on } \mathcal{H} \otimes L^2(\mathbb{T}^n).$$

It has been proved in [6, 11, 15, 16, 21, 29, 41, 42] that the Floquet operator K_F is of pure point spectra or no absolutely continuous spectra where $\mathcal{B}$ is bounded. When $\mathcal{B}$ is unbounded, the first result was obtained by Bambusi and Graffi [1] where they considered the time dependent Schrödinger equation

$$i\partial_t \psi(x, t) = H(t) \psi(x, t), \quad x \in \mathbb{R}; \quad H(t) := -\frac{d^2}{dx^2} + Q(x) + \varepsilon V(x, \omega t), \quad \varepsilon \in \mathbb{R},$$

where $Q(x) \sim |x|^{2\alpha}$ with $\alpha > 1$ as $|x| \rightarrow \infty$ and $|V(x, \theta)| |x|^{-\beta}$ is bounded as $|x| \rightarrow \infty$ for some $\beta < \alpha - 1$. This entails the pure-point nature of the spectrum of the Floquet operator

$$K_F := -i\omega \cdot \partial_\theta - \frac{d^2}{dx^2} + Q(x) + \varepsilon V(x, \theta),$$

on $L^2(\mathbb{R}) \otimes L^2(\mathbb{T}^n)$ for ε small. Liu and Yuan [38] solved the case when $\beta \leq \alpha - 1$. Very recently Bambusi [2, 3] solved the case when $\beta < \alpha + 1$ under some additional assumptions.

For 1-d quantum harmonic oscillator the main difficulty encountered by the traditional KAM method seems to be the eigenvalue spacing for the unperturbed operator does not grow. In [20] Enss and Veselic proved that, if ω is rational, the Floquet operator relative with the 1-d quantum harmonic oscillator has pure point spectrum when the perturbing potential V is bounded and has sufficiently fast decay at infinity. In [13] Combes obtained the reducibility under time periodic, spatially localized perturbation. In [48] Wang proved the spectrum of the Floquet operator K is pure point for the quasiperiodic case where the perturbing potential has *exponential decay*. Grebért and Thomann [24] improved the results in [48] from exponential decay to *polynomial decay*. In [49] we extended the results in [24] from polynomial decay to *logarithmic decay*. Quite recently, in [2, 3] Bambusi dealt with the unbounded perturbation case for 1d harmonic oscillators. For example he can deal with the case $-\partial_{xx} + x^2 + \varepsilon x a_1(\omega t) - i a_2(\omega t) \varepsilon \partial_x$. As Bambusi [2] pointed it out that his results didn't contradict with the interesting counterexamples in [14] and [21].

The results about the reducibility for higher spatial dimension are very few. In [19] Eliasson and Kuksin obtained the reducibility for the Schrödinger equation on $\mathbb{T}^d$. In [22] Grébert and Paturel firstly obtained the reducibility for any dimensional harmonic oscillator on $\mathbb{R}^d$ under the temporal quasiperiodic and analytic perturbation. In this paper we will generalize the results in [22] from temporal analytic perturbations to differential perturbations.

Very recently, Bambusi, Grebért, Maspero and Robert [5] proved a reducibility result for a quantum harmonic oscillator in any dimension perturbed by a linear operator which is a polynomial of degree two in $x_j, -i\partial_j$ with coefficients being real analytic in $\theta \in \mathbb{T}^n$. The proof depends on the following key fact: for polynomial Hamiltonians of degree at most 2 the correspondence between classical and quantum mechanics is exact (see also [28]). But the reducibility problem keeps open for the quantum oscillator in arbitrary dimension with more general unbounded perturbations (see [5], page 2).

1.3. Brief description of the setting and main ideas of the proof. We use the notations introduced in [22]. In phase space $(u, \bar{u}) \in \mathcal{H}^0 \times \mathcal{H}^0$ endowed with the symplectic structure $idu \wedge d\bar{u}$, equation (1.1) is Hamiltonian with

$$H = h(u, \bar{u}) + \varepsilon p(\omega t, u, \bar{u}), \quad (1.4)$$

where

$$h(u, \bar{u}) = \int_{\mathbb{R}^d} (u_x \bar{u}_x + |x|^2 u \bar{u}) dx, \quad p(\omega t, u, \bar{u}) = \int_{\mathbb{R}^d} V(\omega t, x) u \bar{u} dx.$$

Expanding u and $\bar{u}$ on the Hermite basis, $u(x) = \sum_{a \in \mathcal{E}} \xi_a \Phi_a(x)$, $\bar{u}(x) = \sum_{a \in \mathcal{E}} \eta_a \Phi_a(x)$, the phase space $(u, \bar{u}) \in \mathcal{H}^0 \times \mathcal{H}^0$ becomes into the phase space $(\xi, \eta) \in Y_0$ (for the definition of Y_s see Subsection 2.1). We endow Y_0 with the symplectic structure $id\xi \wedge d\eta$. In this setting, (1.4) reads as

$$H(t, \xi, \eta) = \sum_{a \in \mathcal{E}} w_a \xi_a \eta_a + \varepsilon p_\omega(t, \xi, \eta) \quad (1.5)$$

where $p_\omega(t, \xi, \eta) = \langle \xi, P(\omega t) \eta \rangle = \sum_{a, b \in \mathcal{E}} P_a^b(\omega t) \xi_a \eta_b$, which is quadratic in (ξ, η) with

$$P_a^b(\omega t) = \int_{\mathbb{R}^d} V(\omega t, x) \Phi_a(x) \Phi_b(x) dx, \quad a, b \in \mathcal{E}. \quad (1.6)$$

Therefore, the reducibility problem of system (1.1) is equivalent to the reducibility problem for the Hamiltonian system

$$\begin{cases} \dot{\xi}_a = -i w_a \xi_a - i \varepsilon (P^T(\omega t) \xi)_a, \\ \dot{\eta}_a = i w_a \eta_a + i \varepsilon (P(\omega t) \eta)_a, \end{cases} \quad a \in \mathcal{E} \quad (1.7)$$

associated to the non autonomous quadratic Hamiltonian function (1.5). We will give a general reducibility result in Subsection 2.2 which can be applied to system (1.7) and the proof is based on KAM theory. We remark that KAM theory is almost well-developed for nonlinear Hamiltonian PDEs in 1-d context. See [4, 26, 30, 31, 32, 33, 34, 35, 36, 37, 44, 51] for 1-d KAM results. Comparing with 1-d case, the KAM results for multidimensional PDEs are relatively few. Refer to [17, 18, 23, 25, 27, 45] for n-d results. See [8] for an almost complete picture of recent KAM theory.

Highlights. By introducing $\theta = \omega t$, system (1.7) is equivalent to an autonomous system with Hamiltonian

$$\mathcal{H}(\theta, y, \xi, \eta) = \sum_{j=1}^n \omega_j y_j + \langle \xi, N_0 \eta \rangle + \varepsilon \langle \xi, P(\theta) \eta \rangle, \quad \langle \xi, N_0 \eta \rangle = \sum_{a \in \mathcal{E}} w_a \xi_a \eta_a. \quad (1.8)$$

In [22] Grébert and Paturel assumed that the potential $V(\theta, \cdot)$ is real analytic with value in $\mathcal{H}^s(\mathbb{R}^d)$ with $s > 2(d-2) \geq 0$ when $d \geq 2$. Here we only discuss the higher dimensional case for simplicity. Then in Lemma 3.2 Grébert and Paturel [22] proved that $P(\theta) \in \mathcal{M}_{s,\alpha}(D_0, \sigma)$, where $\alpha > 0$ is critical (see the definition of $\mathcal{M}_{s,\alpha}(D_0, \sigma)$ in Section 2). In this paper we only assume that the potential $V(\theta, \cdot) \in \mathcal{C}^\beta(\mathbb{T}^n, \mathcal{H}^s(\mathbb{R}^d))$ with the same condition on s . Using the techniques from [22] and functional analysis ([7]) we can prove that $P(\theta) \in \mathcal{C}^\beta(\mathbb{T}^n, \mathcal{M}_{s,\alpha})$ (see Lemma 3.3), which can be considered as a parallel lemma as Lemma 3.2 in [22].

Now our main problem is to build a similar reducibility result for the Hamiltonian (1.8) when $P(\theta) \in \mathcal{C}^\beta(\mathbb{T}^n, \mathcal{M}_{s,\alpha})$ and thus a smooth KAM is needed here. We recall that the smoothing techniques were firstly introduced by Moser [39, 40] and developed later by many people, see Salamon and Zehnder [47], Pöschel [43], Chierchia and Qian [12], Berti and Bolle [9, 10] and etc. An earlier reducibility result about time dependent Schrödinger operator with finite differentiable unbounded perturbation has been obtained by Yuan and Zhang in [50]. In [2] Bambusi's method in dealing with the differential perturbations is more close to the classical proof in [46]. A significant difference between our paper and [50], [2] is that we deal with the matrix block, not the single matrix element. For the following proof we almost follow the presentation of [12] in the spirit of [46] combined with the KAM method in [22].

More clearly, we will introduce a series of analytic functions $P^{(\nu)}(\theta) \in \mathcal{M}_{s,\alpha}(D_0, \sigma_\nu)$, $\nu = 0, 1, 2, \dots$, and $P^{(\nu)}(\theta) \rightarrow P(\theta)$ in $\theta \in \mathbb{T}^n$ as σ_ν shrinking to 0 ($\nu \rightarrow \infty$), see Lemmas 4.7 and 4.8 for details.

Thus, instead of considering the original function $\mathcal{H}$, in each KAM step, we consider the analytic Hamiltonian function

$$H^{(\nu)}(\theta, y, \xi, \eta) = \sum_{j=1}^n \omega_j y_j + \langle \xi, N_0 \eta \rangle + \varepsilon \langle \xi, P^{(\nu)}(\theta) \eta \rangle$$

which is an approximation of (1.8). We suppose that there exists symplectic map Φ^ν such that

$$H^{(\nu)} \circ \Phi^\nu = \sum_{j=1}^n \omega_j y_j + \langle \xi, N_\nu \eta \rangle + \langle \xi, P_\nu(\theta) \eta \rangle$$

with the norm of $P_\nu(\theta)$ is less than $\epsilon_\nu/2$. Then in $(\nu+1)^{th}$ step, we consider the Hamiltonian

$$\begin{aligned} H^{(\nu+1)}(\theta, y, \xi, \eta) &= \sum_{j=1}^n \omega_j y_j + \langle \xi, N_0 \eta \rangle + \varepsilon \langle \xi, P^{(\nu+1)}(\theta) \eta \rangle \\ &= H^{(\nu)} + (H^{(\nu+1)} - H^{(\nu)}) \end{aligned}$$

which is tiny different from $H^{(\nu)}$. By Φ^ν we have

$$H^{(\nu+1)} \circ \Phi^\nu = H^{(\nu)} \circ \Phi^\nu + (H^{(\nu+1)} - H^{(\nu)}) \circ \Phi^\nu.$$

We shrink the radius of the analytic domain from σ_ν to $\sigma_{\nu+1} = \sigma_\nu^{\frac{3}{2}}$ in order to prove that the norm of additional quadratic perturbation term $(H^{(\nu+1)} - H^{(\nu)}) \circ \Phi^\nu$ is less than $\epsilon_\nu/2$ too (see Lemma 4.12). Then from Proposition 4.1([22]) we can construct $\Phi_{\nu+1}$ such that

$$H^{(\nu+1)} \circ \Phi^{\nu+1} = H^{(\nu+1)} \circ \Phi^\nu \circ \Phi_{\nu+1} = h_{\nu+1} + \langle \xi, P_{\nu+1}(\theta) \eta \rangle,$$

where $h_{\nu+1}$ is in normal form and the norm of $P_{\nu+1}$ is less than $\epsilon_{\nu+1}/2$. Thus we can formulate the iteration lemma. Finally, let $\nu \rightarrow \infty$, we obtain $\mathcal{H} \circ \Phi^\infty = \sum_{j=1}^n \omega_j y_j + \langle \xi, N_\infty \eta \rangle$, where $(\xi, \eta) \in Y_{s'}$ with $1 \leq s' \leq \max\{s, 1\}$ and $N_\infty(\omega) = \lim_{\nu \rightarrow \infty} N_\nu(\omega)$ in normal form and with the norm close to N_0 . For obtaining the above proof we need to show that $H^{(\nu)} \rightarrow \mathcal{H}$ and $\Phi^\nu \rightarrow \Phi^\infty$. Furthermore, from the estimation of $\Phi^\nu - \Phi^{\nu-1}$ and Lemma 4.2 we deduce that $\Phi^\infty - id \in \mathcal{C}^\mu(\mathbb{T}^n, \mathfrak{L}(Y_{s'}, Y_{s'+2\alpha}))$ for all $0 \leq s' \leq s$, where $\mu \leq \frac{2}{9}\beta$ and is not an integer.

The paper is organized as follows: In Sect. 2 we state the abstract reducibility theorem: Theorem 2.6. In Sect. 3 we prove Theorem 1.3, Corollary 1.5 and Corollary 1.6, which are direct results from Theorem 2.6. In Sect. 4 we prove Theorem 2.6. The section is split into a few subsections. Finally, the appendix contains some technical lemmas.

2. REDUCIBILITY THEOREM FOR QUANTUM HARMONIC OSCILLATOR IN $\mathbb{R}^d$ WITH QUASIPERIODIC IN TIME POTENTIAL: SMOOTH VERSION.

2.1. Setting. Notations. Denote $\mathbb{C}, \mathbb{R}, \mathbb{Z}, \mathbb{N}$ be the set of all complex numbers, real numbers, integers and nonnegative integers, respectively. $\mathbb{T} = \mathbb{R}/2\pi\mathbb{Z}$. $\langle \cdot, \cdot \rangle$ is the standard scalar product in ℓ^2 , while $\langle f \rangle := \frac{1}{(2\pi)^n} \int_{\mathbb{T}^n} f(\theta) d\theta$ be the mean value of f on the torus $\mathbb{T}^n$. $|\cdot|$ will be general to denote a supremum norm with a notable exception: for a multi-index $k = (k_1, \dots, k_n) \in \mathbb{Z}^n$, denote $|k| = \sum_{i=1}^n |k_i|$. In the whole paper we use ν to stand for the KAM iteration step.

Linear space. Following the notations in Subsection 1.1, for $s \geq 0$, we consider the complex weighted ℓ^2 -space

$$\ell_s^2 := \{ \xi = (\xi_a \in \mathbb{C})_{a \in \mathcal{E}} \mid \|\xi\|_s < \infty \}$$

with $\|\xi\|_s^2 := \sum_{a \in \mathcal{E}} |\xi_a|^2 w_a^s$ and ℓ_0^2 is ℓ^2 . Then we define

$$Y_s := \ell_s^2 \times \ell_s^2 = \{ \zeta = (\zeta_a = (\xi_a, \eta_a) \in \mathbb{C}^2)_{a \in \mathcal{E}} \mid \|\zeta\|_s < \infty \}$$

with $\|\zeta\|_s^2 := \sum_{a \in \mathcal{E}} (|\xi_a|^2 + |\eta_a|^2) w_a^s$.

We provide the space Y_s , $s \geq 0$, with the symplectic structure $i \sum_{a \in \mathcal{E}} d\xi_a \wedge d\eta_a$. To any smooth function $f(\xi, \eta)$ defined on a domain of Y_s , it corresponds to the Hamiltonian system:

$$\begin{cases} \dot{\xi} = -i \nabla_\eta f(\xi, \eta) \\ \dot{\eta} = i \nabla_\xi f(\xi, \eta) \end{cases} \quad (2.1)$$

where $\nabla f = (\nabla_\xi f, \nabla_\eta f)^T$ is the gradient with respect to the scalar product in Y_0 .

For any smooth functions $f(\xi, \eta)$, $g(\xi, \eta)$, the Poisson bracket of f , g is given by

$$\{f, g\} := i \sum_{a \in \mathcal{E}} \left(\frac{\partial f}{\partial \xi_a} \frac{\partial g}{\partial \eta_a} - \frac{\partial g}{\partial \xi_a} \frac{\partial f}{\partial \eta_a} \right). \quad (2.2)$$

We also consider the extended phase space $\mathcal{P}_s := \mathbb{T}^n \times \mathbb{R}^n \times Y_s \ni (\theta, y, \xi, \eta)$. For smooth functions $f(\theta, y, \xi, \eta)$, $g(\theta, y, \xi, \eta)$, the Poisson bracket is given by

$$\{f, g\} := \sum_{j=1}^n \left(\frac{\partial f}{\partial y_j} \frac{\partial g}{\partial \theta_j} - \frac{\partial g}{\partial y_j} \frac{\partial f}{\partial \theta_j} \right) + i \sum_{a \in \mathcal{E}} \left(\frac{\partial f}{\partial \xi_a} \frac{\partial g}{\partial \eta_a} - \frac{\partial g}{\partial \xi_a} \frac{\partial f}{\partial \eta_a} \right). \quad (2.3)$$

Infinite matrices. We denote by $\mathcal{M}_{s,\alpha}$ the set of infinite matrices $A : \mathcal{E} \times \mathcal{E} \rightarrow \mathbb{C}$ with the norm

$$|A|_{s,\alpha} := \sup_{a,b \in \mathcal{E}} (w_a w_b)^\alpha \left\| A_{[a]}^{[b]} \right\| \left(\frac{\sqrt{\min(w_a, w_b)} + |w_a - w_b|}{\sqrt{\min(w_a, w_b)}} \right)^{s/2} < +\infty,$$

where $A_{[a]}^{[b]}$ denotes the restriction of A to the block $[a] \times [b]$ and $\|\cdot\|$ denotes the operator norm.

We also denote $\mathcal{M}_{s,\alpha}^+$ be the subspace of $\mathcal{M}_{s,\alpha}$ satisfying that an infinite matrix $A \in \mathcal{M}_{s,\alpha}^+$ if

$$|A|_{s,\alpha+} := \sup_{a,b \in \mathcal{E}} (w_a w_b)^\alpha (1 + |w_a - w_b|) \left\| A_{[a]}^{[b]} \right\| \left(\frac{\sqrt{\min(w_a, w_b)} + |w_a - w_b|}{\sqrt{\min(w_a, w_b)}} \right)^{s/2} < +\infty.$$

From the definition we have following simple facts.

Lemma 2.1. $(\mathcal{M}_{s,\alpha}, |\cdot|_{s,\alpha})$ and $(\mathcal{M}_{s,\alpha}^+, |\cdot|_{s,\alpha+})$ are Banach spaces.

Lemma 2.2 ([22], Lemma 2.1). *Let $0 < \alpha \leq 1$ and $s \geq 0$, there exists a constant $c(\alpha, s) > 0$ such that*

i) *If $A \in \mathcal{M}_{s,\alpha}$ and $B \in \mathcal{M}_{s,\alpha}^+$, then AB and BA belong to $\mathcal{M}_{s,\alpha}$ and*

$$|AB|_{s,\alpha}, |BA|_{s,\alpha} \leq c(\alpha, s) |A|_{s,\alpha} |B|_{s,\alpha+}.$$

ii) *If $A, B \in \mathcal{M}_{s,\alpha}^+$, then AB belongs to $\mathcal{M}_{s,\alpha}^+$ and*

$$|AB|_{s,\alpha+} \leq c(\alpha, s) |A|_{s,\alpha+} |B|_{s,\alpha+}.$$

iii) *If $A \in \mathcal{M}_{s,\alpha}$, then for any $t \geq 1$, $A \in \mathfrak{L}(\ell_t^2, \ell_{-t}^2)$ and*

$$\|A\xi\|_{-t} \leq c(\alpha, s) |A|_{s,\alpha} \|\xi\|_t.$$

iv) *If $A \in \mathcal{M}_{s,\alpha}^+$, then $A \in \mathfrak{L}(\ell_{s'}^2, \ell_{s'+2\alpha}^2)$ for all $0 \leq s' \leq s$ and*

$$\|A\xi\|_{s'+2\alpha} \leq c(\alpha, s) |A|_{s,\alpha+} \|\xi\|_{s'};$$

furthermore, $A \in \mathfrak{L}(\ell_1^2, \ell_1^2)$ and

$$\|A\xi\|_1 \leq c(\alpha, s) |A|_{s,\alpha+} \|\xi\|_1.$$

Normal form. We introduce the following definitions.

Definition 2.3. A matrix $F : \mathcal{E} \times \mathcal{E} \rightarrow \mathbb{C}$ is Hermitian, i.e., $F_b^a = \overline{F_a^b}$, $a, b \in \mathcal{E}$.

Definition 2.4. A matrix $N : \mathcal{E} \times \mathcal{E} \rightarrow \mathbb{C}$ is in normal form, and we denote $N \in \mathcal{NF}$, if

(i) N is Hermitian,

(ii) N is block diagonal, i.e., $N_b^a = 0$, for $w_a \neq w_b$.

Quadratic form. To a matrix $Q = (Q_a^b)_{a,b \in \mathcal{E}} \in \mathfrak{L}(\ell_t^2, \ell_{-t}^2)$ we associate in a unique way a quadratic form $q(\xi, \eta)$ on Y_t by the formula $q(\xi, \eta) := \langle \xi, Q\eta \rangle = \sum_{a,b \in \mathcal{E}} Q_a^b \xi_a \eta_b$ and the Poisson bracket

$$\{q_1, q_2\}(\xi, \eta) = -i \langle \xi, [Q_1, Q_2] \eta \rangle$$

where $[Q_1, Q_2] = Q_1 Q_2 - Q_2 Q_1$ is the commutator of two matrices Q_1 and Q_2 . Moreover, if $Q \in \mathcal{M}_{s,\alpha}$ then

$$\sup_{a,b \in \mathcal{E}} \left\| (\nabla_\xi \nabla_\eta q)_{[a]}^{[b]} \right\| \leq \frac{|Q|_{s,\alpha}}{(w_a w_b)^\alpha} \left(\frac{\sqrt{\min(w_a, w_b)}}{\sqrt{\min(w_a, w_b)} + |w_a - w_b|} \right)^{s/2}.$$

Parameter. In the paper ω will play the role of a parameter belonging to $D_0 = [0, 2\pi]^n$. All the constructed functions will depend on ω with $\mathcal{C}^1$ regularity. When a function is only defined on a Cantor subset of D_0 the regularity is understood in Whitney sense.

A class of quadratic Hamiltonians. Let $D \subset D_0$, $s \geq 0$, $\alpha > 0$ and $\sigma > 0$. We denote by $\mathcal{M}_{s,\alpha}(D, \sigma)$ the set of mappings as $\mathbb{T}_\sigma^n \times D \ni (\theta, \omega) \mapsto Q(\theta, \omega) \in \mathcal{M}_{s,\alpha}$ which is real analytic on $\theta \in \mathbb{T}_\sigma^n := \{\theta \in \mathbb{C}^n \mid |\Im \theta| < \sigma\}$ and $\mathcal{C}^1$ continuous on $\omega \in D$. This space is equipped with the norm

$$[Q]_{s,\alpha}^{D,\sigma} := \sup_{\omega \in D, |\Im \theta| < \sigma, |k|=0,1} \left| \partial_\omega^k Q(\theta, \omega) \right|_{s,\alpha}.$$

In view of Lemma 2.2, to a matrix $Q \in \mathcal{M}_{s,\alpha}(D, \sigma)$, we can associate the quadratic form on Y_t with $t \geq 1$

$$q(\xi, \eta; \omega, \theta) = \langle \xi, Q(\omega, \theta) \eta \rangle$$

and we have

$$|q(\xi, \eta; \omega, \theta)| \leq c(\alpha, s) [Q]_{s,\alpha}^{D,\sigma} \|(\xi, \eta)\|_t^2$$

for $(\xi, \eta) \in Y_t$, $\omega \in D$, $\theta \in \mathbb{T}_\sigma^n$.

The subspace of $\mathcal{M}_{s,\alpha}(D, \sigma)$ formed by $F(\theta, \omega)$ such that $\partial_\omega^k F(\theta, \omega) \in \mathcal{M}_{s,\alpha}^+$, $|k| = 0, 1$, is denoted by $\mathcal{M}_{s,\alpha}^+(D, \sigma)$ and equipped with the norm

$$[F]_{s,\alpha+}^{D,\sigma} := \sup_{\omega \in D, |\Im \theta| < \sigma, |k|=0,1} \left| \partial_\omega^k F(\theta, \omega) \right|_{s,\alpha+}.$$

The subspace of $\mathcal{M}_{s,\alpha}(D, \sigma)$ that are independent of θ will be denoted by $\mathcal{M}_{s,\alpha}(D)$ and for $N \in \mathcal{M}_{s,\alpha}(D)$,

$$[N]_{s,\alpha}^D := \sup_{\omega \in D, |k|=0,1} \left| \partial_\omega^k N(\omega) \right|_{s,\alpha}.$$

From Lemma 2.2, the following results hold.

Lemma 2.5. For $0 < \alpha \leq 1$ and $s \geq 0$, there exists a constant $c(\alpha, s) > 0$ such that

i) If $A \in \mathcal{M}_{s,\alpha}(D, \sigma)$ and $B \in \mathcal{M}_{s,\alpha}^+(D, \sigma)$, then AB and BA belong to $\mathcal{M}_{s,\alpha}(D, \sigma)$ and

$$[AB]_{s,\alpha}^{D,\sigma}, [BA]_{s,\alpha}^{D,\sigma} \leq c(\alpha, s) [A]_{s,\alpha}^{D,\sigma} [B]_{s,\alpha+}^{D,\sigma}.$$

ii) If $A, B \in \mathcal{M}_{s,\alpha}^+(D, \sigma)$, then AB belongs to $\mathcal{M}_{s,\alpha}^+(D, \sigma)$ and

$$[AB]_{s,\alpha+}^{D,\sigma} \leq c(\alpha, s) [A]_{s,\alpha+}^{D,\sigma} [B]_{s,\alpha+}^{D,\sigma}.$$

Moreover, if $A \in \mathcal{M}_{s,\alpha}(D, \sigma)$ and $B \in \mathcal{M}_{s,\alpha}^+(D, \sigma)$ then Ae^B , $e^B A \in \mathcal{M}_{s,\alpha}(D, \sigma)$ and $e^B - Id \in \mathcal{M}_{s,\alpha}^+(D, \sigma)$ satisfying

$$[Ae^B]_{s,\alpha}, [e^B A]_{s,\alpha} \leq [A]_{s,\alpha}^{D,\sigma} e^{c(\alpha,s)} [B]_{s,\alpha+}^{D,\sigma}. \quad (2.4)$$

$$[e^B - Id]_{s,\alpha+}^{D,\sigma} \leq [B]_{s,\alpha+}^{D,\sigma} e^{c(\alpha,s)} [B]_{s,\alpha+}^{D,\sigma}. \quad (2.5)$$

Hamiltonian flow. When F depends smoothly on θ , $\mathbb{T}^n \ni \theta \mapsto F(\theta) \in \mathcal{M}_{s,\alpha}^+$ with $0 < \alpha \leq 1$ we associate to $f = \langle \xi, F(\theta)\eta \rangle$ the symplectic transformation, generated by the time 1 map of X_f , on the extended phase space $\mathcal{P}_s$:

$$(\theta, y, \xi, \eta) \mapsto (\theta, \tilde{y}, e^{-iF^T} \xi, e^{iF} \eta),$$

where $\tilde{y} = y + \langle \xi, \nabla_\theta F(\theta)\eta \rangle$. In the following we will never calculate $\tilde{y}$ explicitly since the non homogeneous Hamiltonian system (1.7) is equivalent to the system (4.1) where the variable conjugated to θ is not concerned. Thus, the above symplectic transformation is rewritten into a symplectic linear change, restricted in on Y_0 , which is given by $(\xi, \eta) \mapsto (e^{-iF^T} \xi, e^{iF} \eta)$. It is well defined and invertible in Y_0 as a consequence of Lemmas 2.2 and 2.5. Recall that a sufficient and necessary condition for this map to preserve the symmetry $\eta = \bar{\xi}$ is $F^T(\theta) = \overline{F}(\theta)$ when $\theta \in \mathbb{T}^n$, i.e., F is a Hermitian matrix.

C^1 norm of operator in ω . Given $(\theta, \omega) \in \mathbb{T}_\sigma^n \times D$, $\Phi(\theta, \omega) \in \mathfrak{L}(Y_s, Y_{s'})$ being C^1 operator with respect to ω in Whitney sense, we define the C^1 norm of $\Phi(\theta, \omega)$ with respect to ω by

$$\|\Phi\|_{\mathfrak{L}(Y_s, Y_{s'})}^* = \sup_{(\theta, \omega) \in \mathbb{T}_\sigma^n \times D, |k|=0,1, \|\zeta\|_s \neq 0} \frac{\|\partial_\omega^k \Phi(\theta, \omega) \zeta\|_{s'}}{\|\zeta\|_s}.$$

2.2. The reducibility theorem. In this subsection we state an abstract reducibility theorem for quadratic t -quasiperiodic Hamiltonian of the form

$$H(t, \xi, \eta) = \langle \xi, N_0 \eta \rangle + \varepsilon \langle \xi, P(\omega t) \eta \rangle, \quad (\xi, \eta) \in Y_0, \quad (2.6)$$

and the associated Hamiltonian system is

$$\begin{cases} \dot{\xi} = -iN_0 \xi - i\varepsilon P^T(\omega t) \xi, \\ \dot{\eta} = iN_0 \eta + i\varepsilon P(\omega t) \eta, \end{cases} \quad (2.7)$$

where $N_0 = \text{diag}\{\lambda_a, a \in \mathcal{E}\}$ satisfying the following assumptions:

Hypothesis A1 - Asymptotics. There exist positive constants c_0, c_1, c_2 such that

$$c_1 w_a \geq \lambda_a \geq c_2 w_a \text{ and } |\lambda_a - \lambda_b| \geq c_0 |w_a - w_b|, \quad a, b \in \mathcal{E}.$$

Hypothesis A2 - Second Melnikov condition in measure estimates. There exist positive constants α_1, α_2 and c_3 such that the following holds: for each $0 < \kappa < 1/4$ and $K > 0$ there exists a closed subset $D' := D'(\kappa, K) \subset D_0$ with $\text{Meas}(D_0 \setminus D') \leq c_3 K^{\alpha_1} \kappa^{\alpha_2}$ such that for all $\omega \in D'$, $k \in \mathbb{Z}^n$ with $0 < |k| \leq K$ and $a, b \in \mathcal{E}$ we have

$$|\langle k, \omega \rangle + \lambda_a - \lambda_b| \geq \kappa(1 + |w_a - w_b|).$$

Then we have the following reducibility results.

Theorem 2.6. *Given a non autonomous Hamiltonian (2.6) with $d \geq 1$, we assume that $(\lambda_a)_{a \in \mathcal{E}}$ satisfies Hypothesis A1-A2 and $P(\theta) \in \mathcal{C}^\beta(\mathbb{T}^n, \mathcal{M}_{s,\alpha})$ with $s \geq 0$, $\alpha > 0$ and $\beta > \max\{9(2 + \frac{d}{\alpha}) \frac{\gamma_1}{\gamma_2 - 24\delta}, 9n, 12(d+1)\}$ where $\gamma_1 = \max\{\alpha_1, n + d + 2\}$, $\gamma_2 = \frac{\alpha\alpha_2}{4+d+2\alpha\alpha_2}$, $\delta \in (0, \frac{\gamma_2}{24})$.*

Then there exists $\varepsilon_*(n, \beta, s, d, \delta) > 0$ such that if $0 < \varepsilon < \varepsilon_*(n, \beta, s, d, \delta)$, there exist

- (i) *a Cantor set $D_\varepsilon \subset D_0$ with $\text{Meas}(D_0 \setminus D_\varepsilon) \leq c(n, \beta, d, s, \delta) \varepsilon^{\frac{3\delta}{2+d/\alpha}}$;*
- (ii) *a C^1 family in $\omega \in D_\varepsilon$ (in Whitney sense), linear, unitary and symplectic coordinate transformation $\Phi_\omega^\infty(\theta) : Y_0 \rightarrow Y_0$, $\theta \in \mathbb{T}^n$, $\omega \in D_\varepsilon$, of the form*

$$(\xi_+, \eta_+) \mapsto (\xi, \eta) = \Phi_\omega^\infty(\theta)(\xi_+, \eta_+) = (\overline{M}_\omega(\theta)\xi_+, M_\omega(\theta)\eta_+),$$

where $\Phi_\omega^\infty(\theta) - \text{id} \in \mathcal{C}^\mu(\mathbb{T}^n, \mathcal{L}(Y_{s'}, Y_{s'+2\alpha}))$ ($0 \leq s' \leq s, \mu \leq \frac{2}{9}\beta, \mu \notin \mathbb{Z}$) and satisfies

$$\|\Phi_\omega^\infty - \text{id}\|_{\mathcal{C}^\mu(\mathbb{T}^n, \mathcal{L}(Y_{s'}, Y_{s'+2\alpha}))} \leq C(n, \beta, \mu, d, s) \varepsilon^{\frac{3}{2\beta}(\frac{2}{9}\beta - \mu)};$$

- (iii) *a C^1 family of autonomous quadratic Hamiltonians in normal forms*

$$H_\infty(\xi_+, \eta_+) = \langle \xi_+, N_\infty(\omega)\eta_+ \rangle, \quad \omega \in D_\varepsilon,$$

where $N_\infty(\omega) \in \mathcal{NF}$, in particular block diagonal (i.e. $N_a^b = 0$ for $w_a \neq w_b$), and is close to N_0 , i.e. $[N_\infty(\omega) - N_0]_{s,\alpha}^{D_\varepsilon} \leq c(n, \beta, d, s)\varepsilon$, such that

$$H(t, \Phi_\omega^\infty(\omega t)(\xi_+, \eta_+)) = H_\infty(\xi_+, \eta_+), \quad t \in \mathbb{R}, \quad (\xi_+, \eta_+) \in Y_{s'}, \quad \omega \in D_\varepsilon,$$

where $1 \leq s' \leq \max\{s, 1\}$.

Remark 2.7. In fact, $\Phi_\omega^\infty(\theta)$ and its inverse are bounded operators from Y_1 into itself for any $s \geq 0$.

We prove Theorem 2.6 in Section 4.

3. APPLICATION TO THE QUANTUM HARMONIC OSCILLATOR—PROOF OF THEOREM 1.3

In this section we prove Theorem 1.3 as a corollary of Theorem 2.6.

3.1. Verification of the hypothesis.

Lemma 3.1. When $\lambda_a = w_a$, $a \in \mathcal{E}$, Hypothesis A1 holds true with $c_0 = c_1 = c_2 = 1$.

As [22],

Lemma 3.2. When $\lambda_a = w_a$, $a \in \mathcal{E}$, Hypothesis A2 holds true with $D_0 = [0, 2\pi]^n$, $\alpha_1 = n+1$, $\alpha_2 = 1$, $c_3 = c(n)$ and

$$D' := \{\omega \in [0, 2\pi]^n \mid |\langle k, \omega \rangle + j| \geq \kappa(1 + |j|), \text{ for all } j \in \mathbb{Z} \text{ and } k \in \mathbb{Z}^n \setminus \{0\}\}.$$

Proof. Since $w_a - w_b \in \mathbb{Z}$, it is obtained by a straightforward computation. $\square$

Lemma 3.3. Let $d \geq 1$. Suppose that the potential $V : \mathbb{T}^n \times \mathbb{R}^d \ni (\theta, x) \rightarrow \mathbb{R}$ is (s, β) -admissible. Then there exists $\alpha = \alpha(d, s) > 0$ (see (3.1)) such that the matrix function $P(\theta)$ defined by

$$(P(\theta))_a^b = \int_{\mathbb{R}^d} V(\theta, x) \Phi_a(x) \Phi_b(x) dx, \quad a, b \in \mathcal{E},$$

belongs to $\mathcal{C}^\beta(\mathbb{T}^n, \mathcal{M}_{s,\alpha})$.

Proof. We divide the proof into several steps.

(a) We show that

$$\sup_{\theta \in \mathbb{R}^n} |P(\theta)|_{s,\alpha} \leq C(d, s) \|V(\theta, \cdot)\|_{\mathcal{C}^\beta(\mathbb{T}^n, \mathcal{H}^s)}.$$

Recall that for $a, b \in \mathcal{E}$,

$$(P(\theta))_a^b = \int_{\mathbb{R}^d} V(\theta, x) \Phi_a(x) \Phi_b(x) dx.$$

To estimate $|P(\theta)|_{s,\alpha}$ by definition, we turn to estimate

$$\|P_{[a]}^{[b]}(\theta)\| = \sup_{\substack{\Psi_a(x) \in E_{w_a}, \Psi_b(x) \in E_{w_b}, \\ \|\Psi_a(x)\|_{L^2(\mathbb{R}^d)} = \|\Psi_b(x)\|_{L^2(\mathbb{R}^d)} = 1}} \int_{\mathbb{R}^d} V(\theta, x) \Psi_a(x) \Psi_b(x) dx.$$

From a similar proof in [22](Lemma 3.2),

$$\begin{aligned} & \left| \int_{\mathbb{R}^d} V(\theta, x) \Psi_a(x) \Psi_b(x) dx \right| \\ & \leq \frac{C(d, s)}{(w_a w_b)^{\tilde{\alpha}(p)/2}} \left(\frac{\sqrt{\min(w_a, w_b)}}{\sqrt{\min(w_a, w_b) + |w_a - w_b|}} \right)^{s/2} \|V(\theta, \cdot)\|_{\mathcal{H}^s(\mathbb{R}^d)} \end{aligned}$$

where $\tilde{\alpha}(p) = \frac{1}{12}$ if $d = 1$ and $\tilde{\alpha}(p) = \frac{1}{3p}$ ($p \geq \frac{10}{3}$) if $d = 2$ and $\tilde{\alpha}(p) = \frac{1}{2} \left(\frac{d}{3p} - \frac{d-2}{6} \right) > 0$ if $d > 2$ and $\frac{2(d+3)}{d+1} < p < \frac{2d}{d-2}$. Set

$$\alpha := \frac{\tilde{\alpha}(p)}{2} > 0. \quad (3.1)$$

It follows for $\theta \in \mathbb{R}^n$,

$$|P(\theta)|_{s, \alpha} \leq C(d, s) \|V(\theta, \cdot)\|_{\mathcal{H}^s} \leq C(d, s) \|V(\theta, \cdot)\|_{\mathcal{C}^\beta(\mathbb{T}^n, \mathcal{H}^s)}.$$

(b) We show that $P(z) \in \mathcal{C}^0(\mathbb{R}^n, \mathcal{M}_{s, \alpha})$. For any $z_1, z_2 \in \mathbb{R}^n$, $a, b \in \mathcal{E}$,

$$(P(z_1) - P(z_2))_a^b = \int_{\mathbb{R}^d} (V(z_1, x) - V(z_2, x)) \Phi_a(x) \Phi_b(x) dx.$$

A similar discussion as above tells us

$$|P(z_1) - P(z_2)|_{s, \alpha} \leq C(d, s) \|V(z_1, \cdot) - V(z_2, \cdot)\|_{\mathcal{H}^s}.$$

It follows that $|P(z_1) - P(z_2)|_{s, \alpha} \rightarrow 0$ as $z_1 \rightarrow z_2$ by $V(z, \cdot) \in \mathcal{C}^\beta(\mathbb{R}^n, \mathcal{H}^s(\mathbb{R}^d))$.

(c) We show that $P(z)$ is Fréchet differentiable at each $z_0 \in \mathbb{R}^n$ and for $h \in \mathbb{R}^n$, $a, b \in \mathcal{E}$,

$$(P'(z_0)h)_a^b = \int_{\mathbb{R}^d} \langle V'_z(z_0, x), h \rangle \Phi_a(x) \Phi_b(x) dx$$

satisfying

$$\|P'(z_0)\|_{\mathcal{L}(\mathbb{R}^n, \mathcal{M}_{s, \alpha})} \leq C(d, s) \|V(z, \cdot)\|_{\mathcal{C}^\beta(\mathbb{R}^n, \mathcal{H}^s)}. \quad (3.2)$$

In fact, we define for $h \in \mathbb{R}^n$, $a, b \in \mathcal{E}$,

$$(\mathcal{A}h)_a^b := \int_{\mathbb{R}^d} \langle V'_z(z_0, x), h \rangle \Phi_a(x) \Phi_b(x) dx.$$

Clearly, $\mathcal{A}$ is a linear map on $\mathbb{R}^n$. Since $\langle V'_z(z_0, \cdot), h \rangle \in \mathcal{H}^s(\mathbb{R}^d)$, from a similar discussion in (a) we obtain

$$|\mathcal{A}h|_{s, \alpha} \leq C(d, s) \|V'_z(z_0, \cdot)\|_{\mathcal{L}(\mathbb{R}^n, \mathcal{H}^s)} \|h\| \leq C(d, s) \|V(\theta, \cdot)\|_{\mathcal{C}^\beta(\mathbb{R}^n, \mathcal{H}^s)} \|h\|.$$

It follows that

$$\|\mathcal{A}\|_{\mathcal{L}(\mathbb{R}^n, \mathcal{M}_{s, \alpha})} \leq C(d, s) \|V(z, \cdot)\|_{\mathcal{C}^\beta(\mathbb{R}^n, \mathcal{H}^s)}.$$

Next we show that

$$|P(z) - P(z_0) - \mathcal{A}(z - z_0)|_{s, \alpha} = o(|z - z_0|), \quad z \rightarrow z_0. \quad (3.3)$$

For $a, b \in \mathcal{E}$,

$$(P(z) - P(z_0) - \mathcal{A}(z - z_0))_a^b = \int_{\mathbb{R}^d} (V(z, x) - V(z_0, x) - V'_z(z_0, x)(z - z_0)) \Phi_a(x) \Phi_b(x) dx.$$

Note $V(z, \cdot) \in \mathcal{C}^\beta(\mathbb{R}^n, \mathcal{H}^s(\mathbb{R}^d))$, it follows from Taylor's theorem

$$\|V(z, \cdot) - V(z_0, \cdot) - V'_z(z_0, \cdot)(z - z_0)\|_{\mathcal{H}^s} = o(|z - z_0|), \quad z \rightarrow z_0. \quad (3.4)$$

From a similar discussion as above we obtain

$$|P(z) - P(z_0) - \mathcal{A}(z - z_0)|_{s, \alpha} \leq C(d, s) \|V(z, \cdot) - V(z_0, \cdot) - V'_z(z_0, \cdot)(z - z_0)\|_{\mathcal{H}^s}.$$

Combining with (3.4) we have (3.3). Thus $P(z)$ is Fréchet differentiable on $z = z_0$ and $P'(z_0) = \mathcal{A}$ which satisfies (3.2).

(d) We show that $P(z) \in \mathcal{C}^1(\mathbb{R}^n, \mathcal{M}_{s,\alpha})$. Note that for any $z_1, z_2 \in \mathbb{R}^n$, $a, b \in \mathcal{E}$,

$$((P'(z_1) - P'(z_2))h)_a^b = \int_{\mathbb{R}^d} \langle V'_z(z_1, x) - V'_z(z_2, x), h \rangle \Phi_a(x) \Phi_b(x) dx.$$

Thus

$$\begin{aligned} |(P'(z_1) - P'(z_2))h|_{s,\alpha} &\leq C(d, s) \|\langle V'_z(z_1, \cdot) - V'_z(z_2, \cdot), h \rangle\|_{\mathcal{H}^s} \\ &\leq C(d, s) \|V'_z(z_1, \cdot) - V'_z(z_2, \cdot)\|_{\mathfrak{L}(\mathbb{R}^n, \mathcal{H}^s)} \|h\|. \end{aligned}$$

It follows that

$$\|P'(z_1) - P'(z_2)\|_{\mathfrak{L}(\mathbb{R}^n, \mathcal{M}_{s,\alpha})} \leq C(d, s) \|V'_z(z_1, \cdot) - V'_z(z_2, \cdot)\|_{\mathfrak{L}(\mathbb{R}^n, \mathcal{H}^s)}.$$

It follows that $\|V'_z(z_1, \cdot) - V'_z(z_2, \cdot)\|_{\mathfrak{L}(\mathbb{R}^n, \mathcal{H}^s)} \rightarrow 0$ as $z_1 \rightarrow z_2$ by $V(z, \cdot) \in \mathcal{C}^\beta(\mathbb{R}^n, \mathcal{H}^s(\mathbb{R}^d))$ which means that $\|P'(z_1) - P'(z_2)\|_{\mathfrak{L}(\mathbb{R}^n, \mathcal{M}_{s,\alpha})} \rightarrow 0$ as $z_1 \rightarrow z_2$.

(e) Inductively, we assume that $P(z) \in \mathcal{C}^m(\mathbb{R}^n, \mathcal{M}_{s,\alpha})$, $m \leq \beta - 1$, with

$$\left(P^{(m)}(z)(h_1, \dots, h_m)\right)_a^b = \int_{\mathbb{R}^d} V_z^{(m)}(z, x)(h_1, \dots, h_m) \Phi_a(x) \Phi_b(x) dx$$

satisfying

$$\|P^{(m)}(z)\|_{\mathfrak{L}_m(\mathbb{R}^n, \mathcal{M}_{s,\alpha})} \leq C(d, s) \|V(z, \cdot)\|_{\mathcal{C}^\beta(\mathbb{R}^n, \mathcal{H}^s)}, \quad (3.5)$$

where $\mathfrak{L}_m(\mathbb{R}^n, \mathcal{M}_{s,\alpha})$ denotes the multi-linear operator space $\mathfrak{L}(\underbrace{\mathbb{R}^n \times \dots \times \mathbb{R}^n}_m, \mathcal{M}_{s,\alpha})$. Then we

show that $P(z) \in \mathcal{C}^{m+1}(\mathbb{R}^n, \mathcal{M}_{s,\alpha})$ with

$$\left(P^{(m+1)}(z)(h_1, \dots, h_{m+1})\right)_a^b = \int_{\mathbb{R}^d} V_z^{(m+1)}(z, x)(h_1, \dots, h_{m+1}) \Phi_a(x) \Phi_b(x) dx$$

and (3.5) holds for m replaced by $m+1$. We follow the method in steps (c) and (d), and divide the proof into two parts (e_1) and (e_2) respectively.

(e_1) We show that $P^{(m)}(z)$ is Fréchet differentiable on $z = z_0$ and for $a, b \in \mathcal{E}$,

$$\left(P^{(m+1)}(z_0)(h_1, \dots, h_{m+1})\right)_a^b = \int_{\mathbb{R}^d} V_z^{(m+1)}(z_0, x)(h_1, \dots, h_{m+1}) \Phi_a(x) \Phi_b(x) dx$$

with

$$\|P^{(m+1)}(z_0)\|_{\mathfrak{L}_{m+1}(\mathbb{R}^n, \mathcal{M}_{s,\alpha})} \leq C(d, s) \|V(z, \cdot)\|_{\mathcal{C}^\beta(\mathbb{R}^n, \mathcal{H}^s)}. \quad (3.6)$$

In fact, given $z \in \mathbb{R}^n$, we define for $h_1, \dots, h_{m+1} \in \mathbb{R}^n$, $a, b \in \mathcal{E}$,

$$(\mathcal{B}(z)(h_1, \dots, h_{m+1}))_a^b := \int_{\mathbb{R}^d} V_z^{(m+1)}(z, x)(h_1, \dots, h_{m+1}) \Phi_a(x) \Phi_b(x) dx.$$

Clearly, $\mathcal{B} \in \mathfrak{L}_{m+1}(\mathbb{R}^n, \mathcal{M}_{s,\alpha})$. Since $V_z^{(m+1)}(z, \cdot) \in \mathfrak{L}_{m+1}(\mathbb{R}^n, \mathcal{H}^s)$, $\mathcal{B}$ is clearly multi-linear and $V_z^{(m+1)}(z, \cdot)(h_1, \dots, h_{m+1}) \in \mathcal{H}^s(\mathbb{R}^d)$. Combining with

$$\sup_{z \in \mathbb{R}^n} \|V_z^{(m+1)}(z)\|_{\mathfrak{L}_{m+1}(\mathbb{R}^n, \mathcal{H}^s)} \leq \|V(z, \cdot)\|_{\mathcal{C}^\beta(\mathbb{R}^n, \mathcal{H}^s)}$$

and a similar discussion as above we obtain

$$\begin{aligned} |\mathcal{B}(z)(h_1, \dots, h_{m+1})|_{s,\alpha} &\leq C(d, s) \|V_z^{(m+1)}(z)(h_1, \dots, h_{m+1})\|_{\mathcal{H}^s} \\ &\leq C(d, s) \|V(z, \cdot)\|_{\mathcal{C}^\beta(\mathbb{R}^n, \mathcal{H}^s)} \|h_1\| \dots \|h_{m+1}\|. \end{aligned}$$

It follows that

$$\|\mathcal{B}\|_{\mathfrak{L}_{m+1}(\mathbb{R}^n, \mathcal{M}_{s,\alpha})} \leq C(d, s) \|V(z, \cdot)\|_{\mathcal{C}^\beta(\mathbb{R}^n, \mathcal{H}^s)}.$$

Set $B = \mathcal{B}(z_0)$. Next we show that

$$|P^{(m)}(z) - P^{(m)}(z_0) - B(z - z_0)|_{s,\alpha} = o(|z - z_0|), \quad z \rightarrow z_0.$$

For $a, b \in \mathcal{E}$,

$$\begin{aligned} & \left(\left(P^{(m)}(z) - P^{(m)}(z_0) - B(z - z_0) \right) (h_1, \dots, h_m) \right)_a^b \\ &= \int_{\mathbb{R}^d} \left(V_z^{(m)}(z, x) - V_z^{(m)}(z_0, x) - V_z^{(m+1)}(z_0, x)(z - z_0) \right) (h_1, \dots, h_m) \Phi_a(x) \Phi_b(x) dx. \end{aligned}$$

Note $V(z, \cdot) \in \mathcal{C}^\beta(\mathbb{R}^n, \mathcal{H}^s(\mathbb{R}^d))$, it follows from Taylor's theorem

$$\|V_z^{(m)}(z, \cdot) - V_z^{(m)}(z_0, \cdot) - V_z^{(m+1)}(z_0, \cdot)(z - z_0)\|_{\mathfrak{L}_{m+1}(\mathbb{R}^n, \mathcal{H}^s)} = o(|z - z_0|), \quad z \rightarrow z_0.$$

From a similar discussion as in (c) we obtain

$$\| \left(P^{(m)}(z) - P^{(m)}(z_0) - B(z - z_0) \right) (h_1, \dots, h_m) \|_{s, \alpha} \leq C(d, s) o(|z - z_0|) \|h_1\| \cdots \|h_m\|, \quad z \rightarrow z_0.$$

Thus $P^{(m)}(z)$ is Fréchet differentiable and $P^{(m+1)}(z) = \mathcal{B}$ which satisfies (3.6).

(e₂) Since

$$\|V_z^{(m+1)}(z_1, \cdot) - V_z^{(m+1)}(z_2, \cdot)\|_{\mathfrak{L}_{m+1}(\mathbb{R}^n, \mathcal{H}^s)} \rightarrow 0 \text{ as } z_1 \rightarrow z_2$$

by $V(z, \cdot) \in \mathcal{C}^\beta(\mathbb{R}^n, \mathcal{H}^s(\mathbb{R}^d))$. From a similar discussion in (d) we have

$$\begin{aligned} & \|P^{(m+1)}(z_1) - P^{(m+1)}(z_2)\|_{\mathfrak{L}_{m+1}(\mathbb{R}^n, \mathcal{M}_{s, \alpha})} \\ & \leq C(d, s) \|V_z^{(m+1)}(z_1, \cdot) - V_z^{(m+1)}(z_2, \cdot)\|_{\mathfrak{L}_{m+1}(\mathbb{R}^n, \mathcal{H}^s)}, \end{aligned} \quad (3.7)$$

which means that $P(z) \in \mathcal{C}^{m+1}(\mathbb{R}^n, \mathcal{M}_{s, \alpha})$ and

$$\|P^{(m+1)}(z)\|_{\mathfrak{L}_{m+1}(\mathbb{R}^n, \mathcal{M}_{s, \alpha})} \leq C(d, s) \|V(z, \cdot)\|_{\mathcal{C}^\beta(\mathbb{R}^n, \mathcal{H}^s)}.$$

(f) Finally, consider $P(z) \in \mathcal{C}^\beta(\mathbb{R}^n, \mathcal{M}_{s, \alpha})$. Denote $b := \beta - [\beta]$, we show that for any $1 \leq l \leq [\beta]$ integer and $0 < |z_1 - z_2| < 2\pi$,

$$\frac{\|P^{(l)}(z_1) - P^{(l)}(z_2)\|_{\mathfrak{L}_l(\mathbb{R}^n, \mathcal{M}_{s, \alpha})}}{|z_1 - z_2|^b} \leq \|V(z, \cdot)\|_{\mathcal{C}^\beta(\mathbb{R}^n, \mathcal{H}^s)}. \quad (3.8)$$

In fact, note that $V(z, \cdot) \in \mathcal{C}^\beta(\mathbb{R}^n, \mathcal{H}^s(\mathbb{R}^d))$, from the definition we obtain that for $0 < |z_1 - z_2| < 2\pi$,

$$\frac{\|V_z^{(l)}(z_1, \cdot) - V_z^{(l)}(z_2, \cdot)\|_{\mathfrak{L}_l(\mathbb{R}^n, \mathcal{M}_{s, \alpha})}}{|z_1 - z_2|^b} \leq \|V(z, \cdot)\|_{\mathcal{C}^\beta(\mathbb{R}^n, \mathcal{H}^s)}.$$

Thus (3.8) holds by (3.7) for any $1 \leq l \leq [\beta]$ which means that $P(z) \in \mathcal{C}^\beta(\mathbb{R}^n, \mathcal{M}_{s, \alpha})$ and

$$\|P(\theta)\|_{\mathcal{C}^\beta(\mathbb{T}^n, \mathcal{M}_{s, \alpha})} \leq C(d, s) \|V(z, \cdot)\|_{\mathcal{C}^\beta(\mathbb{R}^n, \mathcal{H}^s)}.$$

□

3.2. Proof of Theorem 1.3. Following the discussions stated in Subsection 1.1, the Schrödinger equation (1.1) is equivalent to Hamiltonian system with (1.4). Expanding it on the Hermite basis $(\Phi_a)_{a \in \mathcal{E}}$, it is equivalent to the system governed by (1.5) which reads as (1.7), a special case of system (2.7) with $\lambda_a = w_a$ and $P(\omega t)$ satisfying (1.6). By Lemmas given above, if V is (s, β) -admissible, we can apply Theorem 2.6 to (1.7) with $\gamma_1 = n + d + 2$, $\gamma_2 = \frac{\alpha}{4+d+2\alpha}$ and α given by (3.1), this leads to Theorem 1.3.

More precisely, in the new coordinates given by Theorem 2.6, $(\xi, \eta) = (\overline{M}_\omega \xi_+, M_\omega \eta_+)$, system (1.7) becomes autonomous and decomposes into blocks as follows:

$$\begin{cases} \dot{\xi}_{+, [a]} = -i(\overline{N}_\infty)_{[a]} \xi_{+, [a]}, \\ \dot{\eta}_{+, [a]} = i(N_\infty)_{[a]} \eta_{+, [a]}, \end{cases} \quad a \in \mathcal{E},$$

where $N_\infty = N_\infty(\omega) \in \mathcal{NF}$. Hence the solution start from $(\xi_+(0), \eta_+(0))$ is given by

$$(\xi_+(t), \eta_+(t)) = (e^{-it\overline{N}_\infty} \xi_+(0), e^{itN_\infty} \eta_+(0)), \quad t \in \mathbb{R}.$$

Then the solution $u(t, x)$ of (1.1) corresponding to the initial data $u_0(x) = \sum_{a \in \mathcal{E}} \xi_a(0) \Phi_a(x) \in \mathcal{H}^{s'}$ with $1 \leq s' \leq \max\{s, 1\}$ is formulated by $u(t, x) = \sum_{a \in \mathcal{E}} \xi_a(t) \Phi_a(x)$ with

$$\xi(t) = \overline{M}_\omega(\omega t) e^{-it\overline{N}_\infty} M_\omega^T(0) \xi(0),$$

where we use the fact $(\overline{M}_\omega)^{-1} = M_\omega^T$.

In other words, let us define the transformation $\Psi_\omega(\theta) \in \mathfrak{L}(\mathcal{H}^{s'})$, $0 \leq s' \leq s$, by

$$\Psi_\omega(\theta) \left(\sum_{a \in \mathcal{E}} \xi_a \Phi_a(x) \right) := \sum_{a \in \mathcal{E}} (M_\omega^T(\theta) \xi)_a \Phi_a(x) = \sum_{a \in \mathcal{E}} \xi_{+,a} \Phi_a(x).$$

From a straightforward computation (the proof is given in the Appendix), we have

Lemma 3.4. *For $0 \leq s' \leq s$ and $\alpha > 0$ given by (3.1),*

$$\|\Psi_\omega(\theta) - id\|_{C^\mu(\mathbb{T}^n, \mathfrak{L}(\mathcal{H}^{s'}, \mathcal{H}^{s'+2\alpha}))} \leq C \varepsilon^{\frac{3}{2\beta}(\frac{2}{9}\beta - \mu)},$$

where μ is defined in Theorem 2.6.

Moreover, $u(t, x)$ satisfies (1.1) if and only if $v(t, x) = \Psi_\omega(\omega t) u(t, x)$ satisfies the autonomous equation:

$$i\partial_t v + (-\partial_{xx} + |x|^2)v + \varepsilon W v = 0,$$

where $W(\sum_{a \in \mathcal{E}} \xi_a \Phi_a(x)) = \frac{1}{\varepsilon} \sum_{a \in \mathcal{E}} ((N_\infty - N_0)^T \xi)_a \Phi_a(x)$. Denote by $(W_a^b)_{a,b \in \mathcal{E}}$ the infinite matrix of the operator W written in the Hermite basis ($W_a^b = \int_{\mathbb{R}^d} W \Phi_a \Phi_b dx$), then W is block diagonal. Denote $\langle V \rangle(x) := \frac{1}{(2\pi)^n} \int_{\mathbb{T}^n} V(\theta, x) d\theta$ the mean value of V on the torus with $(\langle V \rangle_a^b)_{a,b \in \mathcal{E}}$ the corresponding infinite matrix where

$$\langle V \rangle_a^b = \int_{\mathbb{R}^d} \langle V \rangle \Phi_a \Phi_b dx = \frac{1}{(2\pi)^n} \int_{\mathbb{T}^n} P_a^b(\theta) d\theta = (P_a^b)^0$$

with $P_a^b(\theta) = \sum_{k \in \mathbb{Z}^n} (P_a^b)^k e^{ik\theta}$. From (4.19) with $\nu = 0$, $(\tilde{N}_0)_{[a]} = \frac{\varepsilon}{(2\pi)^n} \int_{\mathbb{T}^n} P_{[a]}^{(1)}(\theta) d\theta$. It holds that

$$\begin{aligned} \|(\tilde{N}_0)_{[a]} - \varepsilon P_{[a]}^0\| &= \frac{\varepsilon}{(2\pi)^n} \left\| \int_{\mathbb{T}^n} (P^{(1)}(\theta) - P(\theta))_{[a]} d\theta \right\| \\ &\leq c \varepsilon w_a^{-2\alpha} [P^{(1)}(\theta) - P(\theta)]_{s,\alpha}^{\mathbb{T}^n, D_\varepsilon} \\ &\leq c \varepsilon w_a^{-2\alpha} \sigma_1^\beta = c w_a^{-2\alpha} \varepsilon \epsilon_1 \end{aligned}$$

by Lemma 4.6. On the other hand, from Lemma 4.22, $[N_\infty - N_1]_{s,\alpha}^{\mathbb{T}^n, D_\varepsilon} \leq 2\epsilon_1$. Thus,

$$w_a^{2\alpha} \|(W - \langle V \rangle)_{[a]}\| \leq \frac{w_a^{2\alpha}}{\varepsilon} \|(N_\infty - N_0 - \tilde{N}_0)_{[a]}\| + \frac{w_a^{2\alpha}}{\varepsilon} \|(\tilde{N}_0 - \varepsilon P^0)_{[a]}\| \leq c \varepsilon^{\frac{1}{2}}.$$

Therefore,

$$\|(W_a^b)_{a,b \in \mathcal{E}} - \Pi(\langle V \rangle_a^b)_{a,b \in \mathcal{E}}\|_{\mathfrak{L}(\ell_{s'}^2)} \leq c \varepsilon^{\frac{1}{2}}$$

for $0 \leq s' \leq s$, where Π is the projection on the diagonal blocks.

The proofs of Corollary 1.5 and 1.6 are similar as [22], we omit it for simplicity.

4. PROOF OF THEOREM 2.6

In this section we will use a universal constant C to simplify the proof, which depends on n, β, d, s and is changing in the context.

The system (2.7) is equivalent to the autonomous system:

$$\begin{cases} \dot{\xi}_a = -i\lambda_a \xi_a - i\varepsilon(P^T(\theta)\xi)_a, \\ \dot{\eta}_a = i\lambda_a \eta_a + i\varepsilon(P(\theta)\eta)_a, \\ \dot{y} = -\varepsilon\langle \xi, \nabla_\theta P(\theta)\eta \rangle, \\ \dot{\theta} = \omega, \end{cases} \quad a \in \mathcal{E} \quad (4.1)$$

with the Hamiltonian

$$\mathcal{H}(\theta, y, \xi, \eta, \omega) = \sum_{j=1}^n \omega_j y_j + \sum_{a \in \mathcal{E}} \lambda_a \xi_a \eta_a + \varepsilon \langle \xi, P(\theta)\eta \rangle \quad (4.2)$$

in the extended phase space $\mathcal{P}_s := \mathbb{T}^n \times \mathbb{R}^n \times Y_s$, and λ_a satisfies Hypothesis A1 – A2.

4.1. Analytic approximation to a $\mathcal{C}^\beta$ smooth function. The $\mathcal{C}^\beta$ smooth Hamiltonian function (4.2) can be approximated by a series of *analytic* Hamiltonians

$$H^{(\nu)}(\theta, y, \xi, \eta, \omega) = \sum_{j=1}^n \omega_j y_j + \sum_{a \in \mathcal{E}} \lambda_a \xi_a \eta_a + \varepsilon \langle \xi, P^{(\nu)}(\theta)\eta \rangle, \quad \nu = 1, 2, \dots,$$

where $P^{(\nu)}(\theta)$ will be given in the following. In order to extend the $\mathcal{C}^\beta$ function to a complex neighborhood of $\mathbb{T}^n$, we need the famous results.

Lemma 4.1. (*Jackson, Moser and Zehnder*) Let X be a complex Banach space and $f \in \mathcal{C}^\beta(\mathbb{R}^n, X)$ for some $\beta > 0$ with finite $\mathcal{C}^\beta(\mathbb{R}^n, X)$ norm. Let ϕ be a radical - symmetric, $\mathcal{C}^\infty$ function, having as support the closure of the unit ball centered at the origin, where ϕ is completely flat and take value 1, let $K = \hat{\phi}$ be its Fourier transform. For all $\sigma > 0$ define

$$S_\sigma f(z, \cdot) = K_\sigma * f = \frac{1}{\sigma^n} \int_{\mathbb{R}^n} K\left(\frac{z-y}{\sigma}\right) f(y, \cdot) dy.$$

Then there exists a constant $C > 0$ depending on β, n and X such that the following holds: for any $\sigma > 0$, the function $f_\sigma(z)$ is a real analytic function from $\mathbb{C}^n$ to X such that if

$$\Delta_\sigma^n := \{z \in \mathbb{C}^n \mid |\Im z_j| \leq \sigma, 1 \leq j \leq n\},$$

then for any $\nu \in \mathbb{N}^n$ such that $|\nu| \leq \beta$ one has

$$\sup_{z \in \Delta_\sigma^n} \left\| \partial^\nu f_\sigma(z) - \sum_{|k| \leq \beta - |\nu|} \frac{\partial^{k+\nu} f(\Re z)}{k!} (i\Im z)^k \right\|_{X_\nu} \leq C \|f\|_{\mathcal{C}^\beta(\mathbb{R}^n, X)} \sigma^{\beta - |\nu|},$$

and for all $0 \leq \sigma_1 \leq \sigma$,

$$\sup_{z \in \Delta_{\sigma_1}^n} \|\partial^\nu f_\sigma(z) - \partial^\nu f_{\sigma_1}(z)\|_{X_\nu} \leq C \|f\|_{\mathcal{C}^\beta(\mathbb{R}^n, X)} \sigma^{\beta - |\nu|}.$$

The function f_σ preserves periodicity (i.e. if f is T -periodic in any of its variables z_j , so is f_σ).

The same theorem was also used in [2], [50], etc.

The converse statement of Lemma 4.1 holds only if μ is not an integer. A classical version of this converse result is due to Bernstein and relates the differentiability properties of a periodic function to quantitative estimates for an approximating sequence of trigonometric polynomials. In the following lemma, we suppose X be a complex Banach space as above.

Lemma 4.2. *Let $\ell \geq 0$ and n be a positive integer. Then there exists a constant $c = c(\ell, n)$ such that if $f : \mathbb{R}^n \rightarrow X$ is the limit of a sequence of real analytic maps $f_\nu(x)$ in the strips $|\Im x| \leq \sigma_\nu := \sigma^{(\frac{3}{2})^\nu}$ with $0 < \sigma \leq \frac{1}{4}$ and*

$$f_0 = 0, \quad |f_\nu - f_{\nu-1}|_X \leq c\sigma_\nu^\ell$$

for $|\Im x| \leq \sigma_\nu$, $\nu = 1, 2, \dots$, then $f \in \mathcal{C}^\mu(\mathbb{R}^n, X)$ for every $\mu \leq \ell$ which is not an integer and

$$|f|_{\mathcal{C}^\mu(\mathbb{R}^n, X)} \leq \frac{4c(\ell, n)}{\iota(1-\iota)} \sigma^{\ell-\mu}, \quad 0 < \iota := \mu - [\mu] < 1.$$

For the proof see section 5.

For our applications we choose $f(z) = P(z)$, $X = (\mathcal{M}_{s,\alpha}, |\cdot|_{s,\alpha})$. From Lemma 4.1 we denote

$$S_\sigma P(z) = \sigma^{-n} \int_{\mathbb{R}^n} K\left(\frac{z-y}{\sigma}\right) P(y) dy, \quad z \in \Delta_\sigma^n. \quad (4.3)$$

From Lemma 4.1 again we have that for any $\sigma > 0$, $S_\sigma P(z)$ is a real analytic function from $\mathbb{C}^n$ to $\mathcal{M}_{s,\alpha}$ such that for any $k \in \mathbb{N}^n$ satisfying $|k| \leq \beta$, one has

$$\sup_{z \in \Delta_\sigma^n} \|\partial^k S_\sigma P(z) - \sum_{|m| \leq \beta - |k|} \frac{\partial^{m+k} P(\Re z)}{m!} (i\Im z)^m\|_{X_k} \leq C \|P(\theta)\|_{\mathcal{C}^\beta(\mathbb{T}^n, \mathcal{M}_{s,\alpha})} \sigma^{\beta - |k|}, \quad (4.4)$$

and for all $0 \leq \sigma' \leq \sigma$,

$$\sup_{z \in \Delta_{\sigma'}^n} \|\partial^k S_\sigma P(z) - \partial^k S_{\sigma'} P(z)\|_{X_k} \leq C \|P(\theta)\|_{\mathcal{C}^\beta(\mathbb{T}^n, \mathcal{M}_{s,\alpha})} \sigma^{\beta - |k|}. \quad (4.5)$$

Remark 4.3. *Since $S_\sigma P(z)$ preserves periodicity, we often write $S_\sigma P(\theta)$ instead of $S_\sigma P(z)$. Recall $P(\theta)$ is Hermitian and, from [46], $K(\mathbb{R}^n) \in \mathbb{R}$, then $S_\sigma P(\theta)$ is also Hermitian when $\theta \in \mathbb{T}^n$ by (4.3).*

Suppose $0 < \dots < \sigma_\nu < \dots < \sigma_1 < \sigma_0$. Then we can construct a series of analytic functions $\{S_{\sigma_\nu} P(\theta), \theta \in \mathbb{T}_{\sigma_\nu}^n\}_{\nu \in \mathbb{N}}$. From (4.5) we have

Lemma 4.4. *For $|\Im \theta| \leq \sigma_\nu$, $\nu = 1, 2, \dots$,*

$$|S_{\sigma_\nu} P(\theta) - S_{\sigma_{\nu-1}} P(\theta)|_{s,\alpha} \leq C \sigma_{\nu-1}^\beta \|P(\theta)\|_{\mathcal{C}^\beta(\mathbb{T}^n, \mathcal{M}_{s,\alpha})} \leq C \sigma_{\nu-1}^\beta.$$

Lemma 4.5. *For $|\Im \theta| \leq \sigma_0 \leq 1$,*

$$|S_{\sigma_0} P(\theta)|_{s,\alpha} \leq C. \quad (4.6)$$

Proof. From (4.4), for $|\Im \theta| \leq \sigma_0 \leq 1$ we have $|S_{\sigma_0} P(\theta) - \sum_{|k| \leq \beta} \frac{\partial^k P(\Re \theta)}{k!} (i\Im \theta)^k|_{s,\alpha} \leq C$. Thus,

$$|\sum_{|k| \leq \beta} \frac{\partial^k P(\Re \theta)}{k!} (i\Im \theta)^k|_{s,\alpha} \leq \sum_{|k| \leq \beta} |\partial^k P(\Re \theta)|_{s,\alpha} |\sigma_0^{|k|}| \leq C. \quad \square$$

Lemma 4.6. *For $\theta \in \mathbb{T}^n$, $|S_{\sigma_\nu} P(\theta) - P(\theta)|_{s,\alpha} \leq C \sigma_\nu^\beta$, $\nu = 0, 1, 2, \dots$.*

Proof. From Lemma 4.1, we have for $|\Im z| \leq \sigma_\nu$,

$$|S_{\sigma_\nu} P(z) - \sum_{|k| \leq \beta} \frac{\partial^k P(\Re z)}{k!} (i\Im z)^k|_{s,\alpha} \leq C \|P(\theta)\|_{\mathcal{C}^\beta(\mathbb{T}^n, \mathcal{M}_{s,\alpha})} \sigma_\nu^\beta \leq C \sigma_\nu^\beta.$$

On the other hand, if $z \in \mathbb{R}^d$, $\sum_{0 < |k| \leq \beta} \frac{\partial^k P(\Re z)}{k!} (i\Im z)^k = 0$. Thus for $\theta \in \mathbb{T}^d$ it follows $|S_{\sigma_\nu} P(\theta) - P(\theta)|_{s,\alpha} \leq C \sigma_\nu^\beta$. $\square$

In the following we will write $P^{(\nu)}(\theta) := S_{\sigma_\nu} P(\theta)$ for simplicity. Combining with all the above lemmas, we have

Lemma 4.7.

$$\begin{aligned} [P(\theta)]_{s,\alpha}^{D_0} &\leq C, & \theta \in \mathbb{T}^n; \\ [P^{(0)}(\theta)]_{s,\alpha}^{D_0,\sigma_0} &\leq C; \\ [P^{(\nu)}(\theta) - P(\theta)]_{s,\alpha}^{D_0} &\leq C\sigma_\nu^\beta, & \theta \in \mathbb{T}^n; \\ [P^{(\nu+1)}(\theta) - P^{(\nu)}(\theta)]_{s,\alpha}^{D_0,\sigma_{\nu+1}} &\leq C\sigma_\nu^\beta. \end{aligned} \quad (4.7)$$

Lemma 4.8. For $\theta \in \mathbb{T}^n$,

$$P(\theta) = P^{(0)}(\theta) + \sum_{\nu=0}^{\infty} (P^{(\nu+1)}(\theta) - P^{(\nu)}(\theta)), \quad (4.8)$$

under the assumption

$$B_1. \quad \sum_{\nu=0}^{\infty} \sigma_\nu^\beta < \infty. \quad (4.9)$$

Proof. From Lemma 4.7 and (4.9),

$$|P^{(0)}(\theta) + \sum_{\nu=0}^{\infty} (P^{(\nu+1)}(\theta) - P^{(\nu)}(\theta))|_{s,\alpha} \leq |P^{(0)}(\theta)|_{s,\alpha} + \sum_{\nu=0}^{\infty} |P^{(\nu+1)}(\theta) - P^{(\nu)}(\theta)|_{s,\alpha} \leq C$$

for any $\theta \in \mathbb{T}^n$. Thus for $\theta \in \mathbb{T}^n$, $P^{(0)}(\theta) + \sum_{\nu=0}^{\infty} (P^{(\nu+1)}(\theta) - P^{(\nu)}(\theta))$ is an element in $(\mathcal{M}_{s,\alpha}, |\cdot|_{s,\alpha})$. From Lemma 4.7, $P(\theta)$ is also an element in $(\mathcal{M}_{s,\alpha}, |\cdot|_{s,\alpha})$ for $\theta \in \mathbb{T}^n$. The equality (4.8) follows from (4.7) and (4.9). $\square$

4.2. Homological equation. The proof of Theorem 2.6 is followed by an iterative KAM procedure where in each step we will consider a homological equation of the form

$$\omega \cdot \nabla_\theta F(\theta, \omega) - i[N(\omega), F(\theta, \omega)] + Q(\theta, \omega) = \text{remainder}$$

with $N(\omega), \omega \in D \subset D_0$, in normal form close to $N_0 = \text{diag}(\lambda_a)_{a \in \mathcal{E}}$ and $Q \in \mathcal{M}_{s,\alpha}(D, \sigma)$. We can construct a solution $F \in \mathcal{M}_{s,\alpha}^+(D', \sigma')$, $0 < \sigma' < \sigma$, $D' \subset D$ as in [22].

Proposition 4.9 ([22], Proposition 4.1). *Let $(\theta, \omega) \in \mathbb{T}_\sigma^n \times D$ with $0 < \sigma \leq 1$, $D \subset D_0$. Suppose $D \ni \omega \mapsto N(\omega) \in \mathcal{NF}$ be a $\mathcal{C}^1$ mapping that satisfies $[N - N_0]_{s,\alpha}^D \leq \frac{c_0}{4}$, $Q \in \mathcal{M}_{s,\alpha}(D, \sigma)$ Hermitian, $0 < \kappa < \frac{c_0}{2}$ and $K \geq 1$. Then for any $0 < \sigma' < \sigma$ there exists a subset $D' = D'(\kappa, K) \subset D$, satisfying $\text{Meas}(D \setminus D') \leq c(n, c_0, \alpha, \alpha_2) K^{\gamma_1} \kappa^{\gamma_2}$ with $\gamma_1 = \max\{d+n+2, \alpha_1\}$, $\gamma_2 = \frac{\alpha\alpha_2}{4+d+2\alpha\alpha_2}$, and there exist $\tilde{N} \in \mathcal{M}_{s,\alpha}(D') \cap \mathcal{NF}$, $F \in \mathcal{M}_{s,\alpha}^+(D', \sigma')$ and $R \in \mathcal{M}_{s,\alpha}(D', \sigma')$, $\mathcal{C}^1$ in ω and analytic in θ , such that*

$$\omega \cdot \nabla_\theta F(\theta, \omega) - i[N(\omega), F(\theta, \omega)] = \tilde{N}(\omega) - Q(\theta, \omega) + R(\theta, \omega)$$

for all $(\theta, \omega) \in \mathbb{T}_{\sigma'}^n \times D'$ and

$$\begin{aligned} [F]_{s,\alpha+}^{D',\sigma'} &\leq \frac{c(n, d, s, \alpha) K^{1+d}}{\kappa^{2+d/\alpha} (\sigma - \sigma')^n} [Q]_{s,\alpha}^{D,\sigma}; \\ [\tilde{N}]_{s,\alpha}^{D'} &\leq [Q]_{s,\alpha}^{D,\sigma}; \\ [R]_{s,\alpha}^{D',\sigma'} &\leq \frac{c(n, d, s, \alpha) K^{1+d/2} e^{-(\sigma - \sigma')K/2}}{\kappa^{1+d/(2\alpha)} (\sigma - \sigma')^n} [Q]_{s,\alpha}^{D,\sigma}. \end{aligned}$$

Moreover, $\tilde{N}$, $F(\theta)$ and $R(\theta)$ are Hermitian when $\theta \in \mathbb{T}^n$.

4.3. The KAM Step. As the KAM proof in [12, 50], we begin with the initial Hamiltonian $H^{(0)} = h + q_0$ with $h = \sum_{j=1}^n \omega_j y_j + \sum_{a \in \mathcal{E}} \lambda_a \xi_a \eta_a := \langle \omega, y \rangle + \langle \xi, N_0 \eta \rangle$, and $q_0 = \langle \xi, Q_0(\theta) \eta \rangle$ with $Q_0 = \varepsilon P^{(0)} \in \mathcal{M}_{s,\alpha}(D_0, \sigma_0)$. For simplicity we set $\sigma_0 = 1$, by Lemma 4.7,

$$[Q_0]_{s,\alpha}^{D_0, \sigma_0} = [\varepsilon P^{(0)}(\theta)]_{s,\alpha}^{D_0, \sigma_0} \leq C\varepsilon := \frac{1}{2}\epsilon_0.$$

In the ν th step of the KAM scheme, we consider the Hamiltonian $H^{(\nu)} = h + q_\nu$ defined on $\mathbb{T}_{\sigma_\nu}^n \times \mathbb{R}^n \times Y_s \times D_0$, where $q_\nu = \langle \xi, Q_\nu(\theta) \eta \rangle$ with $Q_\nu = \varepsilon P^{(\nu)}$ analytic in $\mathbb{T}_{\sigma_\nu}^n$.

We set $\Phi^\nu = \Phi^{\nu-1} \circ \Phi_\nu : \mathbb{T}_{\sigma_{\nu+1}}^n \times \mathbb{R}^n \times Y_s \rightarrow \mathbb{T}_{\sigma_0}^n \times \mathbb{R}^n \times Y_s$ for $\omega \in D_\nu$ with $\Phi^0 = id$ and $\Phi_j(\theta, y, \xi, \eta) = X_{f_j}^1(\theta, y, \xi, \eta) = (\theta, \tilde{y}, e^{-iF_j^T(\theta)}\xi, e^{iF_j(\theta)}\eta)$ which is generated by the time 1 map of Hamiltonian function $f_j = \langle \xi, F_j(\theta) \eta \rangle$, $j = 1, \dots, \nu$. Under Φ^j , we suppose that

$$H^{(j)} \circ \Phi^j = (h + q_j) \circ \Phi^j = h_j + p_j, \quad j = 1, \dots, \nu,$$

where $h_j(\theta, y, \xi, \eta, \omega) = \langle \omega, y \rangle + \langle \xi, N_j(\omega) \eta \rangle$, $p_j(\theta, y, \xi, \eta, \omega) = \langle \xi, P_j(\theta, \omega) \eta \rangle$ with $(\theta, y, \xi, \eta, \omega) \in \mathbb{T}_{\sigma_{j+1}}^n \times \mathbb{R}^n \times Y_s \times D_j$, and for $j = 1, \dots, \nu$, the following estimates hold

$$[P_j]_{s,\alpha}^{D_j, \sigma_{j+1}} \leq \frac{1}{2}\epsilon_j; \quad (4.10)$$

$$[F_j]_{s,\alpha+}^{D_j, \sigma_{j+1}} \leq c(n, d, s, \alpha)\epsilon_{j-1}^{\frac{13}{24}}; \quad (4.11)$$

$$[N_j - N_{j-1}]_{s,\alpha}^{D_{j-1}, \sigma_j} \leq \epsilon_{j-1}; \quad (4.12)$$

$$\|\Phi_j - id\|_{\mathfrak{L}(Y_{s'}, Y_{s'+2\alpha})}^* \leq c(n, d, s, \alpha)\epsilon_{j-1}^{\frac{13}{24}}, \quad 0 \leq s' \leq s; \quad (4.13)$$

$$\text{Meas}(D_{j-1} \setminus D_j) \leq cK_{j-1}^{\gamma_1} \kappa_{j-1}^{\gamma_2}. \quad (4.14)$$

In the $(\nu + 1)$ th step we consider

$$H^{(\nu+1)}(\theta, y, \xi, \eta, \omega) = h + q_{\nu+1} = H^{(\nu)} + (q_{\nu+1} - q_\nu)$$

with $(\theta, y, \xi, \eta, \omega) \in \mathbb{T}_{\sigma_{\nu+1}}^n \times \mathbb{R}^n \times Y_s \times D_\nu$. By Φ^ν we have

$$H^{(\nu+1)} \circ \Phi^\nu = H^{(\nu)} \circ \Phi^\nu + (q_{\nu+1} - q_\nu) \circ \Phi^\nu := h_\nu + \tilde{p}_\nu, \quad (4.15)$$

where $\tilde{p}_\nu = p_\nu + (q_{\nu+1} - q_\nu) \circ \Phi^\nu := \langle \xi, \tilde{P}_\nu(\theta) \eta \rangle$.

We make some assumptions on parameters during the KAM iteration. For any $\nu = 0, 1, 2, \dots$, the following assumptions hold:

$$\text{B2.} \quad \sigma_{\nu+1} \leq \frac{1}{2}\sigma_\nu.$$

$$\text{B3.} \quad \sigma_\nu = \epsilon_{\nu-1}^{t_1}, \kappa_\nu = \epsilon_\nu^{t_2}, \quad K_{\nu-1}^{d+1} \leq \epsilon_{\nu-1}^{-\frac{1}{8}} \text{ with } nt_1 \leq \frac{1}{6}, \quad (2 + d/\alpha)t_2 \leq \frac{1}{6}.$$

$$\text{B4.} \quad \epsilon_{\nu+1} = \epsilon_\nu^{\frac{3}{2}}.$$

$$\text{B5.} \quad \beta \geq \frac{3}{2t_1}.$$

$$\text{B6.} \quad e^{-\frac{1}{4}K_\nu\sigma_{\nu+1}} \leq \epsilon_\nu.$$

The explicit expressions of these parameters are given in Subsection 4.4. Under these assumptions, we have

Lemma 4.10. *Under Assumptions B2 - B5, if $c(n, \beta, d, s)(1 + c(\alpha, s)\epsilon_0^{1/2})^2\varepsilon \leq \frac{1}{2}$, then $\tilde{P}_\nu$ is Hermitian when $\theta \in \mathbb{T}^n$ and $[\tilde{P}_\nu]_{s,\alpha}^{D_\nu, \sigma_{\nu+1}} \leq \epsilon_\nu$.*

We need the following two preparation lemmas for Lemma 4.10.

Denote $B_\nu = e^{iF_1} \dots e^{iF_\nu}$, where B_ν is defined on $(\theta, \omega) \in \mathbb{T}_{\sigma_{\nu+1}}^n \times D_\nu$. Note that $F_\nu, \nu = 1, \dots$, are Hermitian matrices, thus

$$B_\nu^{-1}(\theta) = \overline{B_\nu}^T(\theta), \quad \theta \in \mathbb{T}^n. \quad (4.16)$$

Lemma 4.11. *If $4\epsilon_0^{1/24} \leq 1$, then $B_\nu - Id$, $B_\nu^{-1} - Id \in \mathcal{M}_{s,\alpha}^+(D_\nu, \sigma_{\nu+1})$ and*

$$[B_\nu - Id]_{s,\alpha+}^{D_\nu, \sigma_{\nu+1}}, [B_\nu^{-1} - Id]_{s,\alpha+}^{D_\nu, \sigma_{\nu+1}} \leq \epsilon_0^{\frac{1}{2}}$$

under Assumptions B2-B4. Moreover, for $0 \leq \nu_1 < \nu$,

$$[B_{\nu_1} - B_\nu]_{s,\alpha+}^{D_\nu, \sigma_{\nu+1}}, [B_{\nu_1}^{-1} - B_\nu^{-1}]_{s,\alpha+}^{D_\nu, \sigma_{\nu+1}} \leq \epsilon_{\nu_1}^{\frac{1}{2}}.$$

Proof. We only give the estimates on $[B_\nu - Id]_{s,\alpha+}^{D_\nu, \sigma_{\nu+1}}$ and $[B_{\nu_1} - B_\nu]_{s,\alpha+}^{D_\nu, \sigma_{\nu+1}}$. From Lemma 2.5 and (4.11), we have

$$[e^{iF_j} - Id]_{s,\alpha+}^{D_\nu, \sigma_{\nu+1}} \leq [F_j]_{s,\alpha+}^{D,\sigma} e^{c(\alpha,s)[F_j]_{s,\alpha+}^{D,\sigma}} \leq 2\epsilon_{j-1}^{\frac{13}{24}}, \quad j = 1, \dots, \nu,$$

where $e^{c(\alpha,s)\epsilon_{j-1}^{\frac{13}{24}}} \leq 2$. It follows that

$$\begin{aligned} & [e^{iF_1} e^{iF_2} - Id]_{s,\alpha+}^{D_\nu, \sigma_{\nu+1}} \\ & \leq [e^{iF_1} - Id]_{s,\alpha+}^{D_\nu, \sigma_{\nu+1}} + [e^{iF_2} - Id]_{s,\alpha+}^{D_\nu, \sigma_{\nu+1}} + [(e^{iF_1} - Id)(e^{iF_2} - Id)]_{s,\alpha+}^{D_\nu, \sigma_{\nu+1}} \\ & \leq 2\epsilon_0^{\frac{13}{24}} + 2\epsilon_1^{\frac{13}{24}} + 4c(\alpha, s)\epsilon_0^{\frac{13}{24}}\epsilon_1^{\frac{13}{24}} \leq 3\epsilon_0^{\frac{13}{24}} + 3\epsilon_1^{\frac{13}{24}}. \end{aligned}$$

By induction, we obtain

$$[e^{iF_1} \dots e^{iF_\nu} - Id]_{s,\alpha+}^{D_\nu, \sigma_{\nu+1}} \leq 3\epsilon_0^{\frac{13}{24}} + 3\epsilon_1^{\frac{13}{24}} + \dots + 3\epsilon_{\nu-1}^{\frac{13}{24}} \leq \epsilon_0^{\frac{1}{2}} \quad (4.17)$$

by Assumption B4.

Following a similar discussion above, we also have for $0 \leq \nu_1 < \nu$,

$$[e^{iF_{\nu_1+1}} \dots e^{iF_\nu} - Id]_{s,\alpha+}^{D_\nu, \sigma_{\nu+1}} \leq C\epsilon_{\nu_1}^{\frac{13}{24}}. \quad (4.18)$$

Note that for $0 \leq \nu_1 < \nu$, $B_{\nu_1} - B_\nu = B_{\nu_1}(Id - e^{iF_{\nu_1+1}} \dots e^{iF_\nu})$. Thus

$$\begin{aligned} & [B_{\nu_1} - B_\nu]_{s,\alpha+}^{D_\nu, \sigma_{\nu+1}} \\ & \leq [(B_{\nu_1} - Id)(Id - e^{iF_{\nu_1+1}} \dots e^{iF_\nu})]_{s,\alpha+}^{D_\nu, \sigma_{\nu+1}} + [Id - e^{iF_{\nu_1+1}} \dots e^{iF_\nu}]_{s,\alpha+}^{D_\nu, \sigma_{\nu+1}} \\ & \leq c(\alpha, s, n)\epsilon_0^{\frac{1}{2}}\epsilon_{\nu_1}^{\frac{13}{24}} + \epsilon_{\nu_1}^{\frac{13}{24}} \leq \epsilon_{\nu_1}^{\frac{1}{2}}. \end{aligned}$$

by (2.4), (2.5), (4.17) and (4.18). $\square$

Lemma 4.12. *If $c(n, \beta, d, s)(1 + c(\alpha, s)\epsilon_0^{1/2})^2 \varepsilon \leq \frac{1}{2}$, then $[B_\nu^{-1}(Q_{\nu+1} - Q_\nu)B_\nu]_{s,\alpha}^{D_\nu, \sigma_{\nu+1}} \leq \frac{1}{2}\epsilon_\nu$ under Assumptions B2-B5.*

Proof. Firstly, we consider $[(Q_{\nu+1} - Q_\nu)B_\nu]_{s,\alpha}^{D_\nu, \sigma_{\nu+1}}$ where

$$[Q_{\nu+1} - Q_\nu]_{s,\alpha}^{D_\nu, \sigma_{\nu+1}} = \varepsilon[P^{(\nu+1)} - P^{(\nu)}]_{s,\alpha}^{D_\nu, \sigma_{\nu+1}} \leq c(n, \beta, d, s)\varepsilon\sigma_\nu^\beta$$

by Lemma 4.7. Then from Lemma 2.2 and Lemma 4.11, it holds that

$$\begin{aligned} & [(Q_{\nu+1} - Q_\nu)B_\nu]_{s,\alpha}^{D_\nu, \sigma_{\nu+1}} \\ & \leq [(Q_{\nu+1} - Q_\nu)(B_\nu - Id)]_{s,\alpha}^{D_\nu, \sigma_{\nu+1}} + [Q_{\nu+1} - Q_\nu]_{s,\alpha}^{D_\nu, \sigma_{\nu+1}} \\ & \leq c(n, \beta, d, s)(1 + c(\alpha, s)\epsilon_0^{1/2})\varepsilon\sigma_\nu^\beta, \end{aligned}$$

and

$$\begin{aligned}
& [B_\nu^{-1}(Q_{\nu+1} - Q_\nu)B_\nu]_{s,\alpha}^{D_\nu, \sigma_{\nu+1}} \\
& \leq (1 + c(\alpha, s)[B_\nu^{-1} - Id]_{s,\alpha+}^{D_\nu, \sigma_{\nu+1}})[(Q_{\nu+1} - Q_\nu)B_\nu]_{s,\alpha}^{D_\nu, \sigma_{\nu+1}} \\
& \leq c(n, \beta, d, s)(1 + c(\alpha, s)\epsilon_0^{1/2})^2 \varepsilon \sigma_\nu^\beta \leq \frac{1}{2} \epsilon_\nu
\end{aligned}$$

under Assumptions B2-B5. $\square$

Note that $\tilde{P}_\nu = P_\nu + B_\nu^{-1}(Q_{\nu+1} - Q_\nu)B_\nu$, then it is easy to check that $\tilde{P}_\nu$ is Hermitian. Lemma 4.10 is obtained immediately from (4.10) and Lemma 4.12. $\square$

Go back to the Hamiltonian (4.15), we write $\tilde{p}_\nu = \Gamma \tilde{p}_\nu + r_\nu$ and $\tilde{P}_\nu = \Gamma \tilde{P}_\nu + R_\nu$ respectively, where

$$\Gamma \tilde{P}_\nu(\theta, \omega) := \sum_{|k| \leq K_\nu} \tilde{P}_\nu^k(\omega) e^{ik \cdot \theta}, \quad R_\nu(\theta, \omega) := \sum_{|k| > K_\nu} \tilde{P}_\nu^k(\omega) e^{ik \cdot \theta},$$

and define $\tilde{N}_\nu(\omega) \in \mathcal{NF}$ satisfying

$$\left(\tilde{N}_\nu(\omega) \right)_{[a]} := \left(\tilde{P}_\nu^0(\omega) \right)_{[a]}. \quad (4.19)$$

In the following we will use $\Phi_{\nu+1} = X_{f_{\nu+1}}^1$ with $f_{\nu+1} = \langle \xi, F_{\nu+1}(\theta)\eta \rangle$ to put $\Gamma \tilde{p}_\nu$ into normal form. Assume $f_{\nu+1} = \langle \xi, F_{\nu+1}(\theta)\eta \rangle$ satisfying

$$\Gamma \tilde{p}_\nu - \langle \xi, \tilde{N}_\nu \eta \rangle + \{h_\nu, f_{\nu+1}\} = 0, \quad (4.20)$$

then under $\Phi_{\nu+1}(\xi, \eta) = X_{f_{\nu+1}}^1(\xi, \eta) = (e^{-iF_{\nu+1}^T \xi}, e^{iF_{\nu+1} \eta})$, we have $H^{(\nu+1)} \circ \Phi^\nu \circ \Phi_{\nu+1} = h_{\nu+1} + p_{\nu+1}$, where $h_{\nu+1} = \langle \omega, y \rangle + \langle \xi, N_{\nu+1} \eta \rangle$ with $N_{\nu+1} = N_\nu + \tilde{N}_\nu$, and $p_{\nu+1} = \langle \xi, P_{\nu+1}(\theta, \omega)\eta \rangle$ with

$$P_{\nu+1} = -i \int_0^1 e^{-itF_{\nu+1}} [t\Gamma \tilde{P}_\nu + (1-t)\tilde{N}_\nu, F_{\nu+1}] e^{itF_{\nu+1}} dt + e^{-iF_{\nu+1}} R_\nu e^{iF_{\nu+1}}. \quad (4.21)$$

Therefore, we obtain

$$H^{(\nu+1)} \circ \Phi^{\nu+1} = h_{\nu+1} + p_{\nu+1} = \langle \omega, y \rangle + \langle \xi, N_{\nu+1}(\omega)\eta \rangle + \langle \xi, P_{\nu+1}(\theta, \omega)\eta \rangle.$$

In the following we will give the explicit estimates on $F_{\nu+1}$, $N_{\nu+1}$, $P_{\nu+1}$ and $\Phi_{\nu+1} - id$.

First of all, (4.20) is equivalent to

$$\omega \cdot \nabla_\theta F_{\nu+1}(\theta) - i[N_\nu, F_{\nu+1}(\theta)] + \Gamma \tilde{P}_\nu(\theta) = \tilde{N}_\nu + R_\nu(\theta). \quad (4.22)$$

From Proposition 4.9 and Assumption B4, we have from (4.22)

$$[N_\nu - N_0]_{s,\alpha}^{D_\nu} \leq \sum_{j=1}^\nu [N_j - N_{j-1}]_{s,\alpha}^{D_\nu} \leq \epsilon_0 + \dots + \epsilon_{\nu-1} \leq 2\epsilon_0 \leq \frac{c_0}{4}. \quad (4.23)$$

It follows that

$$\|\partial_\omega^j (N_\nu - N_0)_{[a]}\| w_a^{2\alpha} \leq 2\epsilon_0 \quad j = 0, 1.$$

From Proposition 4.9, together with (4.23), Lemma 4.10 and Assumptions B2-B6, if $\kappa_\nu \leq \frac{c_0}{2}$ and $K_\nu \geq 1$ then there exists a subset $D_{\nu+1} \subset D_\nu$ with $\text{Meas}(D_\nu \setminus D_{\nu+1}) \leq cK_\nu^{\gamma_1} \kappa_\nu^{\gamma_2}$, and there exist

$\tilde{N}_\nu \in \mathcal{M}_{s,\alpha}(D_{\nu+1}) \cap \mathcal{NF}$, $F_{\nu+1} \in \mathcal{M}_{s,\alpha}^+(D_{\nu+1}, \sigma_{\nu+2})$, $R_\nu \in \mathcal{M}_\alpha(D_{\nu+1}, \sigma_{\nu+2})$, $\mathcal{C}^1$ in ω and analytic in θ , such that (4.22) holds for all $(\theta, \omega) \in \mathbb{T}_{\sigma_{\nu+2}}^n \times D_{\nu+1}$ and

$$[F_{\nu+1}]_{s,\alpha+}^{D_{\nu+1},\sigma_{\nu+2}} \leq \frac{c(n,d,s,\alpha)\epsilon_\nu K_\nu^{d+1}}{\kappa_\nu^{2+d/\alpha}(\sigma_{\nu+1}-\sigma_{\nu+2})^n} \leq c(n,d,s,\alpha)\epsilon_\nu^{\frac{13}{24}}; \quad (4.24)$$

$$[\tilde{N}_\nu]_{s,\alpha}^{D_{\nu+1}} \leq [\tilde{P}_\nu]_{s,\alpha}^{D_{\nu+1},\sigma_{\nu+1}} \leq \epsilon_\nu. \quad (4.25)$$

$$[R_\nu]_{s,\alpha}^{D_{\nu+1},\sigma_{\nu+2}} \leq \frac{c(n,d,s,\alpha)\epsilon_\nu K_\nu^{1+d/2} e^{-(\sigma_{\nu+1}-\sigma_{\nu+2})K_\nu/2}}{\kappa_\nu^{1+d/2\alpha}(\sigma_{\nu+1}-\sigma_{\nu+2})^n} \leq \epsilon_{\nu+1}. \quad (4.26)$$

$\tilde{N}_\nu$, $F_{\nu+1}(\theta)$ and $R_\nu(\theta)$ are Hermitian when $\theta \in \mathbb{T}^n$. Moreover, from Lemmas 2.2, 2.5 and (4.24),

Lemma 4.13. *If $4c(\alpha,s)\varepsilon^{\frac{1}{24}} \leq 1$, then the symplectic map $\Phi_{\nu+1}(\theta, \omega)$ defined in $\mathbb{T}_{\sigma_{\nu+2}}^n \times D_{\nu+1}$ satisfies*

$$\|\Phi_{\nu+1} - id\|_{\mathfrak{L}(Y_{s'}, Y_{s'+2\alpha})}^* \leq 4c(\alpha,s)\epsilon_\nu^{\frac{13}{24}} \leq \epsilon_\nu^{\frac{1}{2}}.$$

for all $0 \leq s' \leq s$.

Proof. Note that $\Phi_{\nu+1}(\xi, \eta) = X_{f_{\nu+1}}^1(\xi, \eta) = (e^{-iF_{\nu+1}^T}\xi, e^{iF_{\nu+1}}\eta)$, where

$$\begin{aligned} \|\partial_\omega^k(e^{iF_{\nu+1}} - Id)\eta\|_{s'+2\alpha} &\leq c(\alpha,s)[e^{iF_{\nu+1}} - Id]_{s,\alpha+}\|\eta\|_{s'} \\ &\leq c(\alpha,s)[F_{\nu+1}]_{s,\alpha+}^{D_{\nu+1},\sigma_{\nu+2}} e^{c(\alpha,s)[F_{\nu+1}]_{s,\alpha+}^{D_{\nu+1},\sigma_{\nu+2}}} \|\eta\|_{s'} \\ &\leq 2c(\alpha,s)\epsilon_\nu^{\frac{13}{24}} \|\eta\|_{s'} \end{aligned}$$

for $k = 0, 1$ and $0 \leq s' \leq s$ by (4.24), Lemmas 2.2 and 2.5. Similarly,

$$\|\partial_\omega^k(e^{-iF_{\nu+1}^T} - Id)\xi\|_{s'+2\alpha} \leq 2c(\alpha,s)\epsilon_\nu^{\frac{13}{24}} \|\xi\|_{s'}$$

for $k = 0, 1$. Therefore, we have $\|\Phi_{\nu+1} - id\|_{\mathfrak{L}(Y_{s'}, Y_{s'+2\alpha})}^* \leq c(\alpha,s,n,d)\epsilon_\nu^{\frac{13}{24}}$. $\square$

Estimates on the new error term. Recall $p_{\nu+1} = \langle \xi, P_{\nu+1}(\theta)\eta \rangle$ with (4.21), we have

Lemma 4.14. *Under Assumptions B2 - B6, if $\varepsilon \leq c^{-1}(n,d,s,\alpha) \ll 1$ then $P_{\nu+1}(\theta)$ is Hermitian when $\theta \in \mathbb{T}^n$ and $[P_{\nu+1}]_{s,\alpha}^{D_{\nu+1},\sigma_{\nu+2}} \leq \frac{1}{2}\epsilon_{\nu+1}$.*

Proof. It is easy to check that $P_{\nu+1}(\theta)$ is Hermitian from (4.21). The estimate is divided into two parts.

1) From (2.4), (4.16), (4.24) and (4.26),

$$\begin{aligned} [e^{-iF_{\nu+1}} R_\nu e^{iF_{\nu+1}}]_{s,\alpha}^{D_{\nu+1},\sigma_{\nu+2}} &\leq e^{4c(\alpha)[F_{\nu+1}]_{s,\alpha+}^{D_{\nu+1},\sigma_{\nu+2}}} [R_\nu]_{s,\alpha}^{D_{\nu+1},\sigma_{\nu+2}} \\ &\leq e^4 [R_\nu]_{s,\alpha}^{D_{\nu+1},\sigma_{\nu+2}} \leq e^4 \epsilon_\nu^{\frac{1}{8}} \epsilon_\nu^{\frac{3}{2}} \leq \frac{1}{4} \epsilon_\nu^{\frac{3}{2}}. \end{aligned}$$

2) For $0 \leq t \leq 1$, note that $\Gamma \tilde{P}_\nu = \tilde{P}_\nu - R_\nu$,

$$\begin{aligned} &[e^{-itF_{\nu+1}} [t\Gamma \tilde{P}_\nu + (1-t)\tilde{N}_\nu, F_{\nu+1}] e^{itF_{\nu+1}}]_{s,\alpha}^{D_{\nu+1},\sigma_{\nu+2}} \\ &\leq e^{4c(\alpha)[F_{\nu+1}]_{s,\alpha+}^{D_{\nu+1},\sigma_{\nu+2}}} [t\Gamma \tilde{P}_\nu + (1-t)\tilde{N}_\nu, F_{\nu+1}]_{s,\alpha}^{D_{\nu+1},\sigma_{\nu+2}} \\ &\leq 2c(\alpha) e^{4c(\alpha)[F_{\nu+1}]_{s,\alpha+}^{D_{\nu+1},\sigma_{\nu+2}}} [F_{\nu+1}]_{s,\alpha+}^{D_{\nu+1},\sigma_{\nu+2}} ([\Gamma \tilde{P}_\nu]_{s,\alpha}^{D_{\nu+1},\sigma_{\nu+2}} + [\tilde{N}_\nu]_{s,\alpha}^{D_{\nu+1}}) \\ &\leq 4c(\alpha) e^4 \epsilon_\nu^{\frac{1}{24}} \epsilon_\nu^{\frac{3}{2}} \leq \frac{1}{4} \epsilon_\nu^{\frac{3}{2}} \end{aligned}$$

by Lemma 2.5, (2.4), (4.16), (4.24), (4.25), (4.26). $\square$

4.4. Iteration Lemma. To iterate the KAM steps infinitely often we choose sequences for the pertinent parameters. Set $\epsilon_0 = 2\epsilon c(n, \beta, d, s)$, $\sigma_0 = 1$ and suppose

$$\epsilon_{\nu+1} = \epsilon_\nu^{\frac{3}{2}}, \sigma_{\nu+1} = \epsilon_\nu^{\frac{3}{2\beta}}, \kappa_\nu = \epsilon_\nu^{\frac{1}{6(2+d/\alpha)}}, K_\nu = 8|\ln \epsilon_\nu| \epsilon_\nu^{-\frac{3}{2\beta}} \quad (4.27)$$

for $\nu \geq 0$. From a straightforward computation in Subsection 4.3 we have

Lemma 4.15 (Iterative Lemma). *Let $0 < \delta < \frac{\gamma_2}{24}$ and*

$$\beta > \beta_* = \max\{9(2 + d/\alpha) \frac{\gamma_1}{\gamma_2 - 24\delta}, 9n, 12(d+1)\} \quad (4.28)$$

with $\gamma_1 = \max\{d + n + 2, \alpha_1\}$, $\gamma_2 = \frac{\alpha\alpha_2}{4+d+2\alpha\alpha_2}$. If $0 < \varepsilon \leq \varepsilon_*(n, d, s, \delta) \ll 1$ then all the iteration series $\epsilon_\nu, \sigma_\nu, \kappa_\nu$ and K_ν satisfy Assumptions B1 – B6, therefore we have the followings:

Suppose that in $\mathbb{T}_{\sigma_{\nu+1}}^n \times \mathbb{R}^n \times Y_s \times D_\nu$,

$$H^{(\nu)} \circ \Phi^\nu(\theta, y, \xi, \eta, \omega) = (h + q_\nu) \circ \Phi^\nu = h_\nu + p_\nu,$$

where $\Phi^\nu = \Phi_1 \circ \Phi_2 \circ \cdots \circ \Phi_\nu$ and $\Phi_j = X_{f_j}^1 : Y_{s'} \rightarrow Y_{s'}$ for all $0 \leq s' \leq s$, $\omega \in D_j$ and $\theta \in \mathbb{T}_{\sigma_{j+1}}$ and satisfies

$$\|\Phi_j - id\|_{\mathfrak{L}(Y_{s'}, Y_{s'+2\alpha})}^* \leq \epsilon_{j-1}^{\frac{1}{2}}, j = 1, \dots, \nu,$$

and $h_\nu = \langle \omega, y \rangle + \langle \xi, N_\nu \eta \rangle$ in normal form and $p_\nu = \langle \xi, P_\nu \eta \rangle$ where $N_\nu - N_0 \in \mathcal{M}_{s,\alpha}(D_\nu)$, $P_\nu \in \mathcal{M}_{s,\alpha}(D_\nu, \sigma_{\nu+1})$ and the following estimates hold:

$$\begin{aligned} \text{Meas}(D_{j-1} \setminus D_j) &\leq cK_{j-1}^{\gamma_1} \kappa_{j-1}^{\gamma_2}; \\ [F_j]_{s,\alpha+}^{D_j, \sigma_{j+1}} &\leq c(n, d, s, \alpha) \epsilon_{j-1}^{\frac{13}{24}}; \\ [N_j - N_{j-1}]_{s,\alpha}^{D_{j-1}, \sigma_j} &\leq \epsilon_{j-1}; \\ [P_j]_{s,\alpha}^{D_j, \sigma_{j+1}} &\leq \frac{1}{2} \epsilon_j. \end{aligned}$$

Then there exist $D_{\nu+1} \subset D_\nu$ and a mapping $\Phi_{\nu+1} = X_{f_{\nu+1}}^1 : Y_{s'} \rightarrow Y_{s'}$ for all $0 \leq s' \leq s$, $\omega \in D_{\nu+1}$, $\theta \in \mathbb{T}_{\sigma_{\nu+2}}$, and

$$H^{(\nu+1)} \circ \Phi^\nu \circ \Phi_{\nu+1} = (h + q_{\nu+1}) \circ \Phi^{\nu+1} = h_{\nu+1} + p_{\nu+1}$$

and $h_{\nu+1} = \langle \omega, y \rangle + \langle \xi, N_{\nu+1} \eta \rangle$ in normal form and $p_{\nu+1} = \langle \xi, P_{\nu+1} \eta \rangle$ where $N_{\nu+1} - N_0 \in \mathcal{M}_{s,\alpha}(D_{\nu+1})$, $P_{\nu+1} \in \mathcal{M}_{s,\alpha}(D_{\nu+1}, \sigma_{\nu+2})$ and the same estimates hold for $j \leq \nu + 1$.

Moreover, since $N_\nu, P_\nu(\theta)$ are Hermitian when $\theta \in \mathbb{T}^n$, so are $N_{\nu+1}, P_{\nu+1}(\theta), F_{\nu+1}(\theta)$. $\square$

4.5. Transition to the limit and the proof of Theorem 2.6. Set $D_\varepsilon = \bigcap_{\nu=0}^\infty D_\nu$. From (4.27), (4.28) and Lemma 4.15 we have

$$\text{Meas}(D_0 \setminus D_\varepsilon) \leq \sum_{\nu=0}^\infty cK_\nu^{\gamma_1} \kappa_\nu^{\gamma_2} \leq c(n, d, s, \beta, \delta) \varepsilon^{\frac{3\delta}{2+d/\alpha}},$$

if $0 < \varepsilon \leq \varepsilon_*(n, d, s, \beta, \delta)$.

In the following we will show that

Lemma 4.16. For $(\omega, \theta) \in D_\varepsilon \times \mathbb{T}^n$, $\{\Phi^\nu - id\}_\nu$ is a Cauchy sequence in $\mathfrak{L}(Y_{s'}, Y_{s'+2\alpha})$ for all $0 \leq s' \leq s$.

To prove this lemma we need to show that

Lemma 4.17. For $0 \leq s' \leq s$, $0 \leq \nu_1 < \nu_2$ and $(\omega, \theta) \in D_\varepsilon \times \mathbb{T}^n$,

$$\|\Phi_{\nu_1+1} \circ \Phi_{\nu_1+2} \circ \cdots \circ \Phi_{\nu_2} - id\|_{\mathfrak{L}(Y_{s'}, Y_{s'+2\alpha})}^* \leq C \epsilon_{\nu_1}^{\frac{1}{2}}.$$

Proof. Similar to the proof of Lemma 4.13, from Lemma 2.2 iv) and (4.18), for all $0 \leq s' \leq s$, $(\theta, \omega) \in \mathbb{T}^n \times D_\varepsilon$ it holds that

$$\begin{aligned} & \|\Phi_{\nu_1+1} \circ \Phi_{\nu_1+2} \circ \cdots \circ \Phi_{\nu_2} - id\|_{\mathfrak{L}(Y_{s'}, Y_{s'+2\alpha})}^* \\ & \leq c(\alpha, s) \max \left\{ [e^{-iF_{\nu_1+1}^T} \cdots e^{-iF_{\nu_2}^T} - Id]_{s, \alpha+}^{D_\varepsilon}, [e^{iF_{\nu_1+1}} \cdots e^{iF_{\nu_2}} - Id]_{s, \alpha+}^{D_\varepsilon} \right\} \leq C\epsilon_{\nu_1}^{1/2}. \end{aligned}$$

□

From Lemma 4.17, let $\nu_1 = 0$ we have

Corollary 4.18. *For all $0 \leq s' \leq s$, $(\omega, \theta) \in D_\varepsilon \times \mathbb{T}^n$,*

$$\|\Phi^\nu - id\|_{\mathfrak{L}(Y_{s'}, Y_{s'+2\alpha})}^* \leq C\epsilon_0^{\frac{1}{2}}.$$

Proof of Lemma 4.16. Recall that

$$\begin{aligned} \Phi^{\nu_1} - \Phi^{\nu_2} &= \Phi^{\nu_1} \circ (\Phi_{\nu_1+1} \circ \Phi_{\nu_1+2} \circ \cdots \circ \Phi_{\nu_2} - id) \\ &= (\Phi^{\nu_1} - id) \circ (\Phi_{\nu_1+1} \circ \Phi_{\nu_1+2} \circ \cdots \circ \Phi_{\nu_2} - id) + \Phi_{\nu_1+1} \circ \Phi_{\nu_1+2} \circ \cdots \circ \Phi_{\nu_2} - id, \end{aligned}$$

Then by Lemmas 4.17 and Corollary 4.18 we have $\|\Phi^{\nu_1} - \Phi^{\nu_2}\|_{\mathfrak{L}(Y_{s'}, Y_{s'+2\alpha})}^* \leq C\epsilon_{\nu_1}^{\frac{1}{2}}.$

We recall that $\Phi^\nu - id$ is a map from $\mathbb{T}_{\sigma_{\nu+1}}^n \times D_\varepsilon$ to an operator in $\mathfrak{L}(Y_{s'}, Y_{s'+2\alpha})$. From Lemmas 2.1 and 4.16 we denote $\Phi_\omega^\infty - id = \lim_{\nu \rightarrow \infty} (\Phi^\nu - id)$ which is a map from $\mathbb{T}^n \times D_\varepsilon$ to $\mathfrak{L}(Y_{s'}, Y_{s'+2\alpha})$. Furthermore,

Lemma 4.19. *For $(\omega, \theta) \in D_\varepsilon \times \mathbb{T}^n$, $\Phi_\omega^\infty - id \in \mathcal{C}^\mu(\mathbb{T}^n, \mathfrak{L}(Y_{s'}, Y_{s'+2\alpha}))$ for all $0 \leq s' \leq s$, where $\mu \leq \frac{2}{9}\beta$ and is not an integer,*

$$\|\Phi_\omega^\infty - id\|_{\mathcal{C}^\mu(\mathbb{T}^n, \mathfrak{L}(Y_{s'}, Y_{s'+2\alpha}))} \leq \frac{c(n, \beta)}{\iota(1-\iota)} \epsilon_0^{\frac{3}{2\beta}(\frac{2}{9}\beta - \mu)}, \quad 0 < \iota := \mu - [\mu] < 1.$$

Proof.

$$\|\Phi^\nu - \Phi^{\nu-1}\|_{\mathfrak{L}(Y_{s'}, Y_{s'+2\alpha})} \leq C\epsilon_{\nu-1}^{\frac{1}{2}} \leq C\sigma_{\nu+1}^{\frac{2\beta}{9}}.$$

From Lemma 4.2, we deduce that $\Phi_\omega^\infty - id \in \mathcal{C}^\mu(\mathbb{T}^n, \mathfrak{L}(Y_{s'}, Y_{s'+2\alpha}))$ for every $\mu \leq \frac{2}{9}\beta$ which is not an integer and,

$$\|\Phi_\omega^\infty - id\|_{\mathcal{C}^\mu(\mathbb{T}^n, \mathfrak{L}(Y_{s'}, Y_{s'+2\alpha}))} \leq \frac{c(n, \beta)}{\iota(1-\iota)} \sigma_1^{\frac{2}{9}\beta - \mu} = \frac{c(n, \beta)}{\iota(1-\iota)} \epsilon_0^{\frac{3}{2\beta}(\frac{2}{9}\beta - \mu)}, \quad 0 < \iota := \mu - [\mu] < 1.$$

□

Remark 4.20. *In fact we can prove that Φ_ω^∞ is also C^1 smooth in ω in Whitney sense and satisfies a similar estimation as above.*

Lemma 4.21. *For any $y \in \mathbb{R}^n$, $(\xi, \eta) \in Y_{s'}$ with $1 \leq s' \leq \max\{s, 1\}$,*

$$\mathcal{H} \circ \Phi_\omega^\infty(\xi, \eta) = h_\infty(\xi, \eta) := \langle \omega, y \rangle + \langle \xi, N_\infty \eta \rangle,$$

uniformly for $\omega \in D_\varepsilon$ and $\theta \in \mathbb{T}^n$, where $N_\infty(\omega) = \lim_{\nu \rightarrow \infty} N_\nu(\omega)$ with $N_\infty(\omega) \in \mathcal{NF}$, uniformly on D_ε , C^1 Whitney smooth and $[N_\infty - N_0]_{s, \alpha}^{D_\varepsilon} \leq 2\epsilon_0$.

We need to prove a series of preparation lemmas.

Lemma 4.22. For any $y \in \mathbb{R}^n$, $(\xi, \eta) \in Y_{\bar{s}}$ and $\bar{s} \geq 1$,

$$\lim_{\nu \rightarrow \infty} (h_\nu + p_\nu) = h_\infty = \langle \omega, y \rangle + \langle \xi, N_\infty \eta \rangle,$$

uniformly on $D_\varepsilon \times \mathbb{T}^n$ and $[N_\infty - N_0]_{s, \alpha}^{D_\varepsilon} \leq 2\epsilon_0$.

Proof. For $\nu_1 < \nu_2$, from Lemma 4.15, $[N_{\nu_1} - N_{\nu_2}]_{s, \alpha}^{D_\varepsilon} \leq 2\epsilon_{\nu_1}$. It follows that $N_\nu - N_0$ is a Cauchy series in the norm $[\cdot]_{s, \alpha}^{D_\varepsilon, \mathbb{T}^n}$ by (4.23). We denote $N_\infty = \lim N_\nu$ by Lemma 2.1. Clearly, $\nu \geq 0$,

$$[N_\infty - N_\nu]_{s, \alpha}^{D_\varepsilon} \leq \sum_{k=\nu}^{\infty} [\tilde{N}_k]_{\alpha}^{D_\varepsilon} \leq 2\epsilon_\nu.$$

Thus

$$|\langle \xi, (N_\infty - N_\nu)\eta \rangle| \leq \|\xi\|_{\bar{s}} \|(N_\infty - N_\nu)\eta\|_{-\bar{s}} \leq c(\alpha, s)[N_\infty - N_\nu]_{s, \alpha}^{D_\varepsilon} \|\xi\|_{\bar{s}} \|\eta\|_{\bar{s}} \leq 2c(\alpha, s)\epsilon_\nu \|\xi\|_{\bar{s}} \|\eta\|_{\bar{s}},$$

which means that $\lim_{\nu} \langle \xi, N_\nu \eta \rangle = \langle \xi, N_\infty \eta \rangle$. On the other hand,

$$|p_\nu| \leq c(\alpha, s)[P_\nu]_{s, \alpha}^{D_\nu, \sigma_{\nu+1}} \|\xi\|_{\bar{s}} \|\eta\|_{\bar{s}} \leq \frac{1}{2}c(\alpha, s)\epsilon_\nu \|\xi\|_{\bar{s}} \|\eta\|_{\bar{s}} \rightarrow 0, \quad \nu \rightarrow \infty.$$

□

Lemma 4.23. If $\epsilon_0 \ll 1$, then for $0 \leq s' \leq s$ and $(\theta, \omega) \in \mathbb{T}^n \times D_\varepsilon$ there exists $M_\omega(\theta)$ defined on $\mathfrak{L}(\ell_{s'}^2, \ell_{s'}^2)(0 \leq s' \leq s)$ such that

- i) $\|B_\nu - M_\omega\|_{\mathfrak{L}(\ell_{s'}^2, \ell_{s'+2\alpha}^2)}, \|B_\nu^{-1} - M_\omega^{-1}\|_{\mathfrak{L}(\ell_{s'}^2, \ell_{s'+2\alpha}^2)} \leq c(\alpha, s, n)\epsilon_\nu^{\frac{1}{2}};$
- ii) $\|M_\omega - Id\|_{\mathfrak{L}(\ell_{s'}^2, \ell_{s'+2\alpha}^2)}, \|M_\omega^{-1} - Id\|_{\mathfrak{L}(\ell_{s'}^2, \ell_{s'+2\alpha}^2)} \leq c(\alpha, s, n)\epsilon_0^{\frac{1}{2}}.$

Proof. From Lemmas 2.2 and 4.11, for $0 \leq \nu_1 < \nu_2$, $\|B_{\nu_1} - B_{\nu_2}\|_{\mathfrak{L}(\ell_{s'}^2, \ell_{s'+2\alpha}^2)} \leq c(\alpha, s, n)\epsilon_{\nu_1}^{\frac{1}{2}}$. It means that $\{B_\nu - Id\}$ is a Cauchy sequence in $\mathfrak{L}(\ell_{s'}^2, \ell_{s'+2\alpha}^2)$ and its limit is denoted by $M_\omega - Id$, which satisfies $\|B_\nu - M_\omega\|_{\mathfrak{L}(\ell_{s'}^2, \ell_{s'+2\alpha}^2)} \leq c(\alpha, s, n)\epsilon_\nu^{\frac{1}{2}}$. Recall that from Lemma 4.11, $[B_\nu - Id]_{s, \alpha+}^{\mathbb{T}^n, D_\varepsilon} \leq \epsilon_0^{\frac{1}{2}}$. It follows that $\|B_\nu - Id\|_{\mathfrak{L}(\ell_{s'}^2, \ell_{s'+2\alpha}^2)} \leq c(\alpha, s, n)\epsilon_0^{\frac{1}{2}}$. Set $\nu \rightarrow \infty$, we obtain

$$\|M_\omega - Id\|_{\mathfrak{L}(\ell_{s'}^2, \ell_{s'+2\alpha}^2)} \leq c(\alpha, s, n)\epsilon_0^{\frac{1}{2}}.$$

The estimates on B_ν^{-1} and M_ω^{-1} are similar, we omit it for simplicity.

□

It is easy to show $M_\omega^{-1}(\theta) = \overline{M_\omega^T}(\theta)$ from (4.16) when $\theta \in \mathbb{T}^n$. Moreover, we can improve Lemma 4.23 to the following

Lemma 4.24. If $\epsilon_0 \ll 1$, then

- i) $[B_\nu - M_\omega]_{s, \alpha+}^{\mathbb{T}^n, D_\varepsilon}, [B_\nu^{-1} - M_\omega^{-1}]_{s, \alpha+}^{\mathbb{T}^n, D_\varepsilon} \leq c(\alpha, s, n)\epsilon_\nu^{\frac{1}{2}};$
- ii) $[M_\omega - Id]_{s, \alpha+}^{\mathbb{T}^n, D_\varepsilon}, [M_\omega^{-1} - Id]_{s, \alpha+}^{\mathbb{T}^n, D_\varepsilon} \leq c(\alpha, s, n)\epsilon_0^{\frac{1}{2}}.$

Remark 4.25. Note that $M_\omega - Id \in \mathcal{M}_{s, \alpha}^+$ and $\alpha > 0$, it follows from Lemma 2.2 that $M_\omega - Id \in \mathfrak{L}(\ell_1^2, \ell_1^2)$ and satisfies

$$\|(M_\omega - Id)\xi\|_1 \leq c(\alpha, s)[M_\omega - Id]_{s, \alpha+}^{\mathbb{T}^n, D_\varepsilon} \|\xi\|_1 \leq c(\alpha, s, n)\epsilon_0^{\frac{1}{2}} \|\xi\|_1,$$

for $(\theta, \omega) \in \mathbb{T}^n \times D_\varepsilon$. Thus for $(\theta, \omega) \in \mathbb{T}^n \times D_\varepsilon$

$$\|M_\omega \xi\|_1 \leq c(\alpha, s, n)\|\xi\|_1. \quad (4.29)$$

Similarly, for $(\theta, \omega) \in \mathbb{T}^n \times D_\varepsilon$,

$$\|M_\omega^{-1} \xi\|_1 \leq c(\alpha, s, n)\|\xi\|_1. \quad (4.30)$$

Denote $\Phi_\omega^\infty(\xi, \eta) = (\overline{M}_\omega \xi, M_\omega \eta)$. From (4.29) and (4.30) Φ_ω^∞ and its inverse are bounded operators from Y_1 into Y_1 .

In the following we denote $U_\nu := B_\nu^{-1} N_0 B_\nu$, $U_\infty := M_\omega^{-1} N_0 M_\omega$, $V_\nu := B_\nu^{-1} P^{(\nu)} B_\nu$ and $V_\infty := \varepsilon M_\omega^{-1} P(\theta) M_\omega$ for simplicity. We will prove that

Lemma 4.26. *For any $y \in \mathbb{R}^n$, $(\xi, \eta) \in Y_{s'}$ with $1 \leq s' \leq \max\{s, 1\}$,*

$$\lim_{\nu \rightarrow \infty} H^{(\nu)} \circ \Phi^\nu(\xi, \eta) = \mathcal{H} \circ \Phi_\omega^\infty(\xi, \eta), \quad (4.31)$$

uniformly for $\omega \in D_\varepsilon$ and $\theta \in \mathbb{T}^n$.

Proof. From the definition, $\Phi^\nu(\xi, \eta) = (\overline{B}_\nu \xi, B_\nu \eta)$ and $(\overline{B}_\nu)^T = B_\nu^{-1}$. Thus,

$$\begin{aligned} H^{(\nu)} \circ \Phi^\nu &= \langle \omega, y \rangle + \langle \overline{B}_\nu \xi, (N_0 + P^{(\nu)}) B_\nu \eta \rangle \\ &= \langle \omega, y \rangle + \langle \xi, B_\nu^{-1} (N_0 + P^{(\nu)}) B_\nu \eta \rangle \\ &= \langle \omega, y \rangle + \langle \xi, U_\nu \eta \rangle + \langle \xi, V_\nu \eta \rangle. \end{aligned}$$

On the other hand, by a straightforward computation, we have

$$\mathcal{H} \circ \Phi_\omega^\infty = \langle \omega, y \rangle + \langle \xi, U_\infty \eta \rangle + \langle \xi, V_\infty \eta \rangle.$$

Then (4.31) is proved by the following two lemmas. □

Lemma 4.27. *For $(\xi, \eta) \in Y_{\bar{s}}$ with $\bar{s} \geq 1$,*

$$\lim_{\nu \rightarrow \infty} \langle \xi, (V_\nu - V_\infty) \eta \rangle = 0,$$

uniformly for $\omega \in D_\varepsilon$, $\theta \in \mathbb{T}^n$.

Proof. Consider

$$\begin{aligned} V_\nu - V_\infty &= B_\nu^{-1} P^{(\nu)} B_\nu - \varepsilon M_\omega^{-1} P M_\omega \\ &= B_\nu^{-1} (P^{(\nu)} - \varepsilon P) B_\nu + \varepsilon (B_\nu^{-1} - M_\omega^{-1}) P B_\nu + \varepsilon B_\nu^{-1} P (B_\nu - M_\omega) \\ &:= I_1 + I_2 + I_3 \end{aligned}$$

We first estimate I_1 . Note for $\theta \in \mathbb{T}^n$, from Lemma 4.8,

$$[P^{(\nu)} - \varepsilon P]_{s, \alpha}^{D_\varepsilon, \mathbb{T}^n} \leq \sum_{m=\nu}^{\infty} [P^{(m+1)} - P^{(m)}]_{s, \alpha}^{D_\varepsilon, \mathbb{T}^n} \leq c(n, \alpha, \beta) \varepsilon \epsilon_\nu.$$

On the other hand, by Lemma 4.11 we have

$$[B_\nu^{-1} - Id]_{s, \alpha+}^{D_\varepsilon, \mathbb{T}^n}, [B_\nu - Id]_{s, \alpha+}^{D_\varepsilon, \mathbb{T}^n} \leq \epsilon_0^{\frac{1}{2}}, \quad (4.32)$$

then, from Lemma 2.5,

$$\begin{aligned} [I_1]_{s, \alpha}^{D_\varepsilon, \mathbb{T}^n} &\leq [(B_\nu^{-1} - Id)(P^{(\nu)} - \varepsilon P)(B_\nu - Id)]_{s, \alpha}^{D_\varepsilon, \mathbb{T}^n} + [(B_\nu^{-1} - Id)(P^{(\nu)} - \varepsilon P)]_{s, \alpha}^{D_\varepsilon, \mathbb{T}^n} \\ &\quad + [(P^{(\nu)} - \varepsilon P)(B_\nu - Id)]_{s, \alpha}^{D_\varepsilon, \mathbb{T}^n} + [P^{(\nu)} - \varepsilon P]_{s, \alpha}^{D_\varepsilon, \mathbb{T}^n} \\ &\leq \epsilon_\nu. \end{aligned}$$

For I_2 , note that $[B_\nu^{-1} - M_\omega^{-1}]_{s, \alpha+}^{D_\varepsilon, \mathbb{T}^n} \leq \epsilon_\nu^{\frac{1}{2}}$ by Lemma 4.24, thus

$$[I_2]_{s, \alpha}^{D_\varepsilon, \mathbb{T}^n} \leq [\varepsilon (B_\nu^{-1} - M_\omega^{-1}) P (B_\nu - Id)]_{s, \alpha}^{D_\varepsilon, \mathbb{T}^n} + [\varepsilon (B_\nu^{-1} - M_\omega^{-1}) P]_{s, \alpha}^{D_\varepsilon, \mathbb{T}^n} \leq c(s, n, \alpha, \beta) \varepsilon \epsilon_\nu^{\frac{1}{2}}$$

combined with (4.32). Similarly, $[I_3]_{s, \alpha}^{D_\varepsilon, \mathbb{T}^n} \leq c(s, n, \alpha, \beta) \varepsilon \epsilon_\nu^{\frac{1}{2}}$. Finally, we have

$$[V_\nu - V_\infty]_{s, \alpha}^{D_\varepsilon, \mathbb{T}^n} \leq c(s, n, \alpha, \beta) \varepsilon \epsilon_\nu^{\frac{1}{2}}, \quad (4.33)$$

and $V_\nu - V_\infty \in \mathfrak{L}(\ell_{\bar{s}}^2, \ell_{-\bar{s}}^2)$. By Lemma 2.5 and (4.33) we obtain

$$|\langle \xi, (V_\nu - V_\infty)\eta \rangle| \leq \|\xi\|_{\bar{s}} \|(V_\nu - V_\infty)\eta\|_{-\bar{s}} \leq c(s, n, \alpha, \beta) \varepsilon \epsilon_\nu^{\frac{1}{2}} \|\xi\|_{\bar{s}} \|\eta\|_{\bar{s}} \rightarrow 0$$

as $\nu \rightarrow 0$ uniformly for $\omega \in D_\varepsilon$, $\theta \in \mathbb{T}^n$.

□

Lemma 4.28. For $(\xi, \eta) \in Y_{s'}$ with $1 \leq s' \leq \max\{s, 1\}$,

$$\lim_{\nu \rightarrow \infty} \langle \xi, (U_\nu - U_\infty)\eta \rangle = 0,$$

uniformly for $\omega \in D_\varepsilon$, $\theta \in \mathbb{T}^n$.

Proof. Consider

$$\begin{aligned} \langle \xi, (U_\nu - U_\infty)\eta \rangle &= \langle \xi, (B_\nu^{-1} N_0 B_\nu - M_\omega^{-1} N_0 M_\omega)\eta \rangle \\ &= \langle \xi, (B_\nu^{-1} - M_\omega^{-1}) N_0 B_\nu \eta \rangle + \langle \xi, M_\omega^{-1} N_0 (M_\omega - B_\nu)\eta \rangle. \end{aligned}$$

We estimate the first term. For $\eta' \in \ell_{s'}^2$, $\|N_0 \eta'\|_{s'-2} \leq c_1 \|\eta'\|_{s'}$ by $\lambda_a \leq c_1 w_a$. From Lemma 4.32 and Lemma 2.2 we obtain $\|N_0 B_\nu \eta\|_{s'-2} \leq c(\alpha, s, n) \|\eta\|_{s'}$. Therefore,

$$\begin{aligned} &|\langle \xi, (B_\nu^{-1} - M_\omega^{-1}) N_0 B_\nu \eta \rangle| \\ &= |\langle (\overline{B}_\nu - \overline{M}_\omega) \xi, N_0 B_\nu \eta \rangle| \\ \text{from Cauchy and } 1 \leq s' \leq \max\{s, 1\} &\leq \|(\overline{B}_\nu - \overline{M}_\omega) \xi\|_{s'} \|N_0 B_\nu \eta\|_{s'-2} \\ \text{Lemma 4.24, Lemma 2.2} &\leq c(\alpha, s, n) \epsilon_\nu^{\frac{1}{2}} \|\xi\|_{s'} \|\eta\|_{s'}. \end{aligned}$$

Similarly, for $(\xi, \eta) \in Y_{s'}$, we have $|\langle \xi, M_\omega^{-1} N_0 (M_\omega - B_\nu)\eta \rangle| \leq c(\alpha, s, n) \epsilon_\nu^{\frac{1}{2}} \|\xi\|_{s'} \|\eta\|_{s'}$. The conclusion is clear now. □

Combining with Lemma 4.22 and Lemma 4.26, we finish the proof of Lemma 4.21. □

From Lemma 4.21 and the concrete form of Φ^∞ , we obtain that

$$(\omega t, \star, \overline{M}_\omega(\omega t) e^{-i\overline{N}_\infty t} \xi_0, M_\omega(\omega t) e^{iN_\infty t} \eta_0)$$

are the solutions of the Hamiltonian system (4.1). Thus, $(\overline{M}_\omega(\omega t) e^{-i\overline{N}_\infty t} \xi_0, M_\omega(\omega t) e^{iN_\infty t} \eta_0)$ are clearly the solutions of the Hamiltonian system (2.7). We complete the proofs of Theorem 2.6. □

5. APPENDIX

5.1. Proof of Lemma 3.4.

Proof. Recall that

$$\Psi_\omega(\theta) \left(\sum_{a \in \mathcal{E}} \xi_a \Phi_a(x) \right) = \sum_{a \in \mathcal{E}} (M_\omega^T(\theta) \xi)_a \Phi_a(x).$$

From Theorem 2.6, it is easy to show by definition $M_\omega^T - Id \in \mathcal{C}^\mu(\mathbb{T}^n, \mathfrak{L}(\ell_{s'}^2, \ell_{s'+2\alpha}^2))$ and

$$\|M_\omega^T - Id\|_{\mathcal{C}^\mu(\mathbb{T}^n, \mathfrak{L}(\ell_{s'}^2, \ell_{s'+2\alpha}^2))} \leq C(n, \beta, \mu, d, s) \epsilon^{\frac{3}{2\beta}(\frac{2}{9}\beta - \mu)}.$$

(a) From definition,

$$\begin{aligned} \|\Psi_\omega(\cdot) - id\|_{\mathfrak{L}(\mathcal{H}^{s'}, \mathcal{H}^{s'+2\alpha})} &= \sup_{\|u\|_{\mathcal{H}^{s'}}=1} \|\Psi_\omega(\cdot)u - u\|_{\mathcal{H}^{s'+2\alpha}} \\ &= \sup_{\|\xi\|_{\ell_{s'}^2}=1} \|M_\omega^T \xi - \xi\|_{\ell_{s'+2\alpha}^2} = \|M_\omega^T - Id\|_{\mathfrak{L}(\ell_{s'}^2, \ell_{s'+2\alpha}^2)} \leq C(n, \beta, \mu, d, s) \epsilon^{\frac{3}{2\beta}(\frac{2}{9}\beta - \mu)}. \end{aligned}$$

(b) For $b = \mu - [\mu] \in (0, 1)$, $z_1, z_2 \in \mathbb{R}^n$ with $0 < |z_1 - z_2| < 2\pi$,

$$\begin{aligned} &\frac{\|\Psi_\omega(z_1) - \Psi_\omega(z_2)\|_{\mathfrak{L}(\mathcal{H}^{s'}, \mathcal{H}^{s'+2\alpha})}}{|z_1 - z_2|^b} \\ &= \frac{1}{|z_1 - z_2|^b} \sup_{\|u\|_{\mathcal{H}^{s'}}=1} \|\Psi_\omega(z_1)u - \Psi_\omega(z_2)u\|_{\mathcal{H}^{s'+2\alpha}} \\ &= \frac{1}{|z_1 - z_2|^b} \sup_{\|\xi\|_{\ell_{s'}^2}=1} \|M_\omega^T(z_1)\xi - M_\omega^T(z_2)\xi\|_{\ell_{s'+2\alpha}^2} \\ &= \frac{1}{|z_1 - z_2|^b} \|M_\omega^T(z_1) - M_\omega^T(z_2)\|_{\mathfrak{L}(\ell_{s'}^2, \ell_{s'+2\alpha}^2)} \end{aligned}$$

which shows that

$$\|\Psi_\omega - id\|_{C^b(\mathbb{T}^n, \mathfrak{L}(\mathcal{H}^{s'}, \mathcal{H}^{s'+2\alpha}))} = \|M_\omega^T - Id\|_{C^b(\mathbb{T}^n, \mathfrak{L}(\ell_{s'}^2, \ell_{s'+2\alpha}^2))} \leq C(n, \beta, \mu, d, s) \epsilon^{\frac{3}{2\beta}(\frac{2}{9}\beta - \mu)}.$$

(c) Denote $\langle \mathcal{A}(z), h \rangle u := \sum_{a \in \mathcal{E}} (\langle A(z), h \rangle \xi)_a \Phi_a(x)$ for $h \in \mathbb{R}^n$ where we use the notation $A := (M_\omega^T - Id)'_z$ and $\xi \in \ell_{s'}^2$ and $0 \leq s' \leq s$. Note $M_\omega^T - Id \in C^\mu(\mathbb{T}^n, \mathfrak{L}(\ell_{s'}^2, \ell_{s'+2\alpha}^2))$, it follows that for any $z \in \mathbb{T}^n$, $\mathcal{A}(z) \in \mathfrak{L}(\mathbb{R}^n, \mathfrak{L}(\mathcal{H}^{s'}, \mathcal{H}^{s'+2\alpha}))$. This is because

$$\begin{aligned} \|\mathcal{A}\|_{\mathfrak{L}(\mathbb{R}^n, \mathfrak{L}(\mathcal{H}^{s'}, \mathcal{H}^{s'+2\alpha}))} &= \sup_{\|u\|_{\mathcal{H}^{s'}}=1, \|h\|=1} \|\langle \mathcal{A}(z), h \rangle u\|_{\mathcal{H}^{s'+2\alpha}} \\ &= \sup_{\|\xi\|_{\ell_{s'}^2}=1, \|h\|=1} \|\langle A(z), h \rangle \xi\|_{\ell_{s'+2\alpha}^2} \\ &= \|A(z)\|_{\mathfrak{L}(\mathbb{R}^n, \mathfrak{L}(\ell_{s'}^2, \ell_{s'+2\alpha}^2))} \\ &\leq \|M_\omega^T - Id\|_{C^b(\mathbb{T}^n, \mathfrak{L}(\ell_{s'}^2, \ell_{s'+2\alpha}^2))}. \end{aligned} \tag{5.1}$$

Given $z_0 \in \mathbb{T}^n$,

$$\begin{aligned} &\|\Psi(z) - \Psi(z_0) - \langle \mathcal{A}(z_0), z - z_0 \rangle\|_{\mathfrak{L}(\mathcal{H}^{s'}, \mathcal{H}^{s'+2\alpha})} \\ &= \sup_{\|u\|_{\mathcal{H}^{s'}}=1} \|(\Psi(z) - \Psi(z_0) - \langle \mathcal{A}(z_0), z - z_0 \rangle)u\|_{\mathcal{H}^{s'+2\alpha}} \\ &= \sup_{\|\xi\|_{\ell_{s'}^2}=1} \|(M_\omega^T(z) - M_\omega^T(z_0) - \langle A(z_0), z - z_0 \rangle)\xi\|_{\ell_{s'+2\alpha}^2} \\ &= \|M_\omega^T(z) - M_\omega^T(z_0) - \langle A(z_0), z - z_0 \rangle\|_{\mathfrak{L}(\ell_{s'}^2, \ell_{s'+2\alpha}^2)} \end{aligned}$$

Note $M_\omega^T - Id \in C^\mu(\mathbb{T}^n, \mathfrak{L}(\ell_{s'}^2, \ell_{s'+2\alpha}^2))$, then

$$\|M_\omega^T(z) - M_\omega^T(z_0) - \langle A(z_0), z - z_0 \rangle\|_{\mathfrak{L}(\ell_{s'}^2, \ell_{s'+2\alpha}^2)} = o(|z - z_0|), \quad z \rightarrow z_0.$$

Therefore,

$$\|\Psi(z) - \Psi(z_0) - \langle \mathcal{A}(z_0), z - z_0 \rangle\|_{\mathfrak{L}(\mathcal{H}^{s'}, \mathcal{H}^{s'+2\alpha})} = o(|z - z_0|), \quad z \rightarrow z_0,$$

which shows that $\Psi(z)$ is Fréchet differentiable at z_0 . Moreover, following (b), we have

$$\begin{aligned}
& \sup_{\substack{\|u\|_{\mathcal{H}^{s'}}=1, \|h\|=1 \\ z_1, z_2 \in \mathbb{R}^n, 0 < |z_1 - z_2| < 2\pi}} \frac{1}{|z_1 - z_2|^b} \|\langle \mathcal{A}(z_1) - \mathcal{A}(z_2), h \rangle u\|_{\mathcal{H}^{s'+2\alpha}} \\
&= \sup_{\substack{\|\xi\|_{\ell^2_{s'}}=1, \|h\|=1 \\ z_1, z_2 \in \mathbb{R}^n, 0 < |z_1 - z_2| < 2\pi}} \frac{1}{|z_1 - z_2|^b} \|\langle A(z_1) - A(z_2), h \rangle \xi\|_{\ell^2_{s'+2\alpha}} \\
&= \sup_{z_1, z_2 \in \mathbb{R}^n, 0 < |z_1 - z_2| < 2\pi} \frac{1}{|z_1 - z_2|^b} \|A(z_1) - A(z_2)\|_{\mathfrak{L}(\mathbb{R}^n, \mathfrak{L}(\ell^2_{s'}, \ell^2_{s'+2\alpha}))}.
\end{aligned}$$

Combining with (5.1), we have

$$\|\Psi_\omega(\cdot) - id\|_{C^{1+b}(\mathbb{T}^n, \mathfrak{L}(\mathcal{H}^{s'}, \mathcal{H}^{s'+2\alpha}))} = \|M_\omega^T(\cdot) - Id\|_{C^{1+b}(\mathbb{T}^n, \mathfrak{L}(\ell^2_{s'}, \ell^2_{s'+2\alpha}))} \leq C(n, \beta, \mu, d, s) \epsilon^{\frac{3}{2\beta}(\frac{2}{9}\beta - \mu)}$$

with $1 + b \leq \mu$. Inductively, we can show that

$$\|\Psi_\omega(\cdot) - id\|_{C^{k+b}(\mathbb{T}^n, \mathfrak{L}(\mathcal{H}^{s'}, \mathcal{H}^{s'+2\alpha}))} = \|M_\omega^T(\cdot) - Id\|_{C^{k+b}(\mathbb{T}^n, \mathfrak{L}(\ell^2_{s'}, \ell^2_{s'+2\alpha}))} \leq C(n, \beta, \mu, d, s) \epsilon^{\frac{3}{2\beta}(\frac{2}{9}\beta - \mu)}$$

with $k + b \leq \mu$. Thus we finish the proof of Lemma 3.4. $\square$

5.2. Proof of Lemma 4.2.

Proof. Following Salamon[46], it is enough to consider the case $\mu = \ell$. Moreover, once the result has been established for $0 < \ell < 1$ it follows for $\ell > 1$ by Cauchy's estimate. Therefore we assume $0 < \iota = \mu = \ell < 1$.

Define $g_\nu = f_\nu - f_{\nu-1}$. Then $f = \sum g_\nu$ satisfies the estimate

$$|f|_X \leq c \sum_{\nu=1}^{\infty} \sigma_\nu^\iota = c \sum_{\nu=1}^{\infty} \sigma^{\iota(\frac{3}{2})^\nu} \leq \frac{2c}{\iota} \sigma^\iota,$$

where we use the fact that $\sum_{\nu=1}^{\infty} \sigma^{\iota(\frac{3}{2})^\nu} \leq \sigma^\iota \sum_{\nu \geq 0} (\frac{1}{2^\nu})^\iota \leq \frac{2}{\iota} \sigma^\iota$ for $0 < \sigma \leq 1/4$.

For $x, y \in \mathbb{R}^n$ with $\sigma < |x - y| \leq 1$ this implies $|f(x) - f(y)|_X \leq \frac{4c}{\iota} \sigma^\iota \leq \frac{4c}{\iota} |x - y|^\iota$. In the case $0 < |x - y| \leq \sigma$ there is an integer $N \geq 0$ such that $\sigma_{N+1} < |x - y| \leq \sigma_N$. Following Cauchy's estimate, $|\partial_x g_\nu(u)|_X \leq c \sigma_\nu^{\iota-1}$ for every $u \in \mathbb{R}^n$, we have $|g_\nu(x) - g_\nu(y)|_X \leq c \sigma_\nu^{\iota-1} |x - y|$. We shall use this estimate for $\nu = 1, 2, \dots, N$. For $\nu \geq N+1$ we use the trivial estimate $|g_\nu(x) - g_\nu(y)|_X \leq 2c \sigma_\nu^\iota$. Taking into account the inequalities we obtain that

$$\begin{aligned}
|f(x) - f(y)|_X &\leq \sum_{1 \leq \nu \leq N} |g_\nu(x) - g_\nu(y)|_X + \sum_{\nu > N} |g_\nu(x) - g_\nu(y)|_X \\
&\leq c|x - y| \sum_{1 \leq \nu \leq N} \sigma_\nu^{\iota-1} + 2c \sum_{\nu > N} \sigma_\nu^\iota \\
&\leq c|x - y| \sigma_N^{\iota-1} \sum_{0 \leq \nu \leq N-1} \left(\frac{1}{2^\nu}\right)^{1-\iota} + 2c \sigma_{N+1}^\iota \sum_{\nu \geq 0} \left(\frac{1}{2^\nu}\right)^\iota \\
&\leq \frac{4c}{1-\iota} |x - y| \sigma_N^{\iota-1} + \frac{4c}{\iota} \sigma_{N+1}^\iota \\
&= \frac{4c}{\iota(1-\iota)} |x - y|^\iota.
\end{aligned}$$

We finish the proof. $\square$

REFERENCES

- [1] Bambusi, D., Graffi, S.: Time quasi-periodic unbounded perturbations of Schrödinger operators and KAM method. *Commun. Math. Phys.* **219**(2), 465-480(2001)
- [2] Bambusi, D.: Reducibility of 1-d Schrödinger equation with time quasiperiodic unbounded perturbations, I. *Trans. AMS*, To appear(2017); arXiv: 1606.04494 [math.DS](2016)
- [3] Bambusi, D.: Reducibility of 1-d Schrödinger equation with time quasiperiodic unbounded perturbations, II. *Commun. Math. Phys.* doi: 10. 1007/s00220-016-2825-2(2017)
- [4] Berti, M., Biasco, L., Procesi, M.: KAM for the reversible derivative wave equation. *Arch. Rational Mech. Anal.* **212**, 905-955(2014)
- [5] Bambusi, D., Grébert, B., Maspero, A., Robert, D.: Reducibility of the quantum harmonic oscillator in d-dimensions with polynomial time dependent perturbation, 1-d Schrödinger equation with time quasiperiodic unbounded perturbations. preprint, arXiv: 1702.05274v1 [math. AP](2017)
- [6] Bellissard, J.: Stability and instability in quantum mechanics. Trends and developments in the eighties(Bielefeld, 1982/1983), 1-106. Singapore: World Scientific, 1985.
- [7] Berger, M.: Nonlinearity and functional analysis. Lectures on nonlinear problems in mathematical analysis. Pure and Applied Mathematics. Academic Press, New York-London, 1977.
- [8] Berti, M.: KAM for PDEs. *Boll. Unione Mat. Ital.* **9**, 115-142(2016)
- [9] Berti, M., Bolle, P.: Sobolev quasiperiodic solutions of multidimensional wave equations with a multiplicative potential. *Nonlinearity* **25**, 2579-2613(2012)
- [10] Berti, M., Bolle, P.: Quasi-periodic solutions with Sobolev regularity of NLS on $\mathbb{T}^d$ with a multiplicative potential. *Eur. J. Math.* **15**, 229-286(2013)
- [11] Blekher, P.M., Jauslin, H.R., Lebowitz, J.L.: Floquet spectrum for two-level systems in quasiperiodic time-dependent fields. *J. Statist. Phys.* **68**(1-2), 271-310(1992)
- [12] Chierchia, L., Qian, D.: Moser's theorem for lower dimensional tori. *J. Diff. Eqs.* **206**, 55-93(2004)
- [13] Combescure, M.: The quantum stability problem for time-periodic perturbations of the harmonic oscillator. *Ann. Inst. H. Poincaré Phys. Théor.* **47**(1), 63-83(1987)
- [14] Delort, J. M.: Growth of Sobolev norms for solutions of time dependent Schrödinger operators with harmonic oscillator potential. *Comm. Partial Differential Equations*, **39**(1), 1-33(2014)
- [15] Duclos, P., Stovicek, P.: Floquet Hamiltonians with pure point spectrum. *Commun. Math. Phys.* **177**, 327-347(1996)
- [16] Duclos, P., Stovicek, P., Vittot, M.: Perturbation of an eigenvalue from a dense point spectrum: A general Floquet Hamiltonian. *An. Inst. H. Poincaré Phys. Théor.* **71**, 241-301(1999)
- [17] Eliasson, H. L., Grébert, B., Kuksin, S. B.: KAM for the nonlinear beam equation. *Geom. Funct. Anal.* **26**(6), 1588-1715(2016)
- [18] Eliasson, H. L., Kuksin, S. B.: KAM for the nonlinear Schrödinger equation. *Ann. of Math.* **172**, 371-435(2010)
- [19] Eliasson, L.H., Kuksin, S.B.: On reducibility of Schrödinger equations with quasiperiodic in time potentials. *Commun. Math. Phys.* **286**(1), 125-135(2009)
- [20] Enss, V., Veselic, K.: Bound states and propagating states for time-dependent Hamiltonians. *Ann IHP* **39**(2), 159-191(1983)
- [21] Graffi, S., Yajima, K.: Absolute continuity of the floquet spectrum for a nonlinearly forced harmonic oscillator. *Commun. Math. Phys.* **215**, 245-250(2000)
- [22] Grébert, B., Paturel, E.: On reducibility of quantum harmonic oscillator on $\mathbb{R}^d$ with quasiperiodic in time potential. preprint, arXiv: 1603.07455v1 [math.AP](2016)
- [23] Grébert, B., Paturel, E.: KAM for the Klein Gordon equation on $\mathbb{S}^d$. *Boll. Unione Mat. Ital.* **9**(2), 237-288(2016)
- [24] Grébert, B., Thomann, L.: KAM for the quantum harmonic oscillator. *Commun. Math. Phys.* **307**, 383-427(2011)
- [25] Geng, J., Xu, X., You, J.: An infinite dimensional KAM theorem and its application to the two dimensional cubic Schrödinger equation. *Adv. Math.* **226**, 5361-5402(2011)
- [26] Geng, J., You, J.: A KAM theorem for one dimensional Schrödinger equation with periodic boundary conditions. *J. Diff. Eqs.* **209**, 1-56(2005)
- [27] Geng, J., You, J.: A KAM theorem for Hamiltonian partial differential equations in higher dimensional spaces. *Commun. Math. Phys.* **262**, 343-372(2006)
- [28] Hagedorn, G., Loss, M., Slawny, J.: Non stochasticity of time-dependent quadratic Hamiltonians and the spectra of canonical transformations. *J. Phys. A* **19**(4), 521-531(1986)
- [29] Howland, J.: Floquet operators with singular spectrum, I. *An. Inst. H. Poincaré* **49**, 309-323(1989)
- [30] Kappeler, T., Liang, Z.: A KAM theorem for the defocusing NLS equation with periodic boundary conditions. *J. Diff. Eqs.* **252**, 4068-4113(2012)
- [31] Kuksin, S.B., Pöschel, J.: Invariant cantor manifolds of quasi-periodic oscillations for a nonlinear Schrödinger equation. *Ann. of Math.* **143**, 149-179(1996)

- [32] Kuksin, S.B.: Nearly integrable infinite-dimensional Hamiltonian systems. Lecture Notes in Mathematics **1556**, Springer-Verlag, Berlin, 1993.
- [33] Kuksin, S.B.: Analysis of Hamiltonina PDEs. Oxford University Press, Oxford, 2000.
- [34] Kuksin, S.B.: A KAM theorem for equations of the Korteweg-de Vries type. Rev. Math-Math Phys. **10**(3), 1-64(1998)
- [35] Liang, Z.: Quasi-periodic solutions for 1D Schrödinger equations with the nonlinearity $|u|^{2p}u$. J. Diff. Eqs. **244**, 2185-2225(2008)
- [36] Liang, Z., You, J.: Quasi-periodic solutions for 1D Schrödinger equations with higher order nonlinearity. SIAM J.Math. Anal. **36**, 1965-1990(2005)
- [37] Liu, J., Yuan, X.: A KAM theorem for Hamiltonian partial differential equations with unbounded perturbations. Comm. Math. Phys. **307**(3), 629-673(2011).
- [38] Liu, J., Yuan, X.: Spectrum for quantum Duffing oscillator and small-divisor equation with large-variable coefficient. Comm. Pure Appl. Math. **63**(9), 1145-1172(2010)
- [39] Moser, J.: A rapidly convergent iteration method and non-linear partial differential equations I and II. Ann Scuola Norm, Sup. Pisa **20**, 265-315, 499-535(1966)
- [40] Moser, J.: On the construction of almost periodic solutions for ordinary differential equations. Proc. Intl. Conf. Funct. Anal. and Rel. Top. Tokyo, 60-67(1969)
- [41] Joye, A.: Absence of absolutely continuous spectrum of Floquet operators. J. Statist. Phys. **75**(5-6), 929-952(1994)
- [42] Nenciu, G.: Floquet operators without absolutely continuous spectrum. Ann. Inst. H. Poincaré **59**, 91-97(1993)
- [43] Pöschel, J.: Invariante tori in differenzierbaren Hamiltonschen systemen. Bonner Mathematische Schriften **120** (1980)
- [44] Pöschel, J.: A KAM theorem for some nonlinear partial differential equations. Ann. Sc. Norm. sup. Pisa Cl. sci. **23**, 119-148(1996)
- [45] Procesi, M., Xu, X.: Quasi-töplitz functions in KAM theorem. SIAM J. Math. Anal. **45**, 2148-2181(2013)
- [46] Salamon, D.A.: The Kolmogorov-Arnold-Moser theorem. Math. Phys. Electron. J., **10**, 3-37(electronic) (2004)
- [47] Salamon, D.A., Zehnder, E.: KAM theory in configuration space. Comment. Math. Helv. **64**, 84-132 (1989)
- [48] Wang, W.: Pure point spectrum of the Floquet Hamiltonian for the quantum harmonic oscillator under time quasi-periodic perturbations. Commun. Math. Phys. **277**(2), 459-496(2008)
- [49] Wang, Z., Liang, Z.: Reducibility of 1d quantum harmonic oscillator perturbed by a quasiperiodic potential with logarithmic decay. Nonlinearity **30**, 1405-1448(2017)
- [50] Yuan, X., Zhang K.: A reduction theorem for time dependent Schrödinger operator with finite differentiable unbounded perturbation. J. Math. Phys. **54**(5), 465-480(2013)
- [51] Zhang, J., Gao, M., Yuan, X.: KAM tori for reversible partial differential equations. Nonlinearity **24**, 1189-1228(2011)

SCHOOL OF MATHEMATICAL SCIENCES AND KEY LAB OF MATHEMATICS FOR NONLINEAR SCIENCE, FUDAN UNIVERSITY, SHANGHAI 200433, CHINA

E-mail address: zgliang@fudan.edu.cn

SCHOOL OF MATHEMATICAL SCIENCES, SOOCHOW UNIVERSITY, SUZHOU 215006, CHINA

E-mail address: zgwang@suda.edu.cn